



# Proteomics assisted profiling of antimicrobial peptide signatures from black pepper (*Piper nigrum* L.)

P. Umadevi<sup>1</sup> · M. Soumya<sup>1</sup> · Johnson K. George<sup>1</sup> · M. Anandaraj<sup>2</sup>

Received: 3 January 2018 / Revised: 7 February 2018 / Accepted: 11 March 2018 / Published online: 3 April 2018  
© Prof. H.S. Srivastava Foundation for Science and Society 2018

**Abstract** Plant antimicrobial peptides are the interesting source of studies in defense response as they are essential components of innate immunity which exert rapid defense response. In spite of abundant reports on the isolation of antimicrobial peptides (AMPs) from many sources, the profile of AMPs expressed/identified from single crop species under certain stress/physiological condition is still unknown. This work describes the AMP signature profile of black pepper and their expression upon *Phytophthora* infection using label-free quantitative proteomics strategy. The differential expression of 24 AMPs suggests that a combinatorial strategy is working in the defense network. The 24 AMP signatures belonged to the cationic, anionic, cysteine-rich and cysteine-free group. As the first report on the possible involvement of AMP signature in *Phytophthora* infection, our results offer a platform for further study on regulation, evolutionary importance and exploitation of these AMPs as next generation molecules against pathogens.

**Keywords** Proteomics · Antimicrobial peptides · Differential expression · Host–pathogen interaction

## Introduction

Antimicrobial peptides (AMP) are small peptides, size ranging from 2 to 9 kDa with broad spectrum antimicrobial activity. The percentage distribution is high in animals (74.53%) followed by plants (13.57%) (Sarika et al. 2012). On the basis of electric charge, the AMPs are either cationic or anionic (Pelegri et al. 2011) and they are constitutively expressed or regulated upon stress (Nawrot et al. 2014). AMPs have been isolated from many plants and from various plant parts viz. leaves, stems, roots, flowers and seeds and were proved to act against phytopathogens. They belong to the families viz., defensins, thionins, lipid transfer protein (LTP), snakings, cyclotides and hevein like proteins which are cysteine rich peptides (Park et al. 2000). Except few reports (Egorov et al. 2005; Silva et al. 2012; Zipfel 2009), the description of cysteine free AMPs from plants are rare. Apart from the direct action against pathogens, plant AMPs are also important molecules in MAPK (MAP Kinase) defense signaling (Scott et al. 2007), innate immunity (Rahnamaeian 2011), ROS and H<sub>2</sub>O<sub>2</sub> accumulation (Fan et al. 2008).

Black pepper is an export oriented spice crop, rich in essential oil and oleoresin. Among the biotic /abiotic stresses, the foot rot disease caused by *Phytophthora* is of major concern (Anandaraj 2000) in black pepper. The investigation on presence of AMPs (both constitutive and induced) and its characterization from the resistant genotype would yield information on innate immunity, which will help in developing resistant varieties and also the candidate AMPs as possible lead molecules in future management strategies.

Chromatography based (Cammue et al. 1992) and EST based (Asiegbu et al. 2003; Ke et al. 2015) methods were used to identify and isolate the AMPs from plants. But the

✉ P. Umadevi  
Umavannan28@gmail.com

<sup>1</sup> Division of Crop Improvement and Biotechnology, ICAR-Indian Institute of Spices Research, Kozhikode, Kerala 673 012, India

<sup>2</sup> Division of Crop Protection, ICAR-Indian Institute of Spices Research, Kozhikode, Kerala 673 012, India

AMPs are underrepresented in these conventional methods, including immunoblots due to their extreme isoelectric points and small size (Zhou et al. 2011). Weinhold et al. 2015 quantified the ectopic expression of AMPs in transgenic *Nicotiana attenuata* from apoplastic proteins using label-free protein quantification by nanoUPLC-MS<sup>E</sup> analysis coupled with Hi3 method. Our present study was aimed to explore the label free proteomics strategy to identify the AMPs in resistant variety of black pepper upon infection by *Phytophthora*. The aim of this work was to bring out the entire profile along with expression quantification of AMP signatures upon infection with *Phytophthora* from the total leaf protein using label free proteomics and in-silico analysis of physiochemical, biological properties of AMP signatures. For the first time, we showed the occurrence of both cysteine rich, non cysteine AMP signatures from a complex sample and some major AMPs as innate immunity factors against *Phytophthora*.

## Materials and methods

### In planta inoculation

Black pepper variety, “IISR Shakthi” resistant to *Phytophthora capsici* was used in this study. The plants with three to four leaves were inoculated at the abaxial side (in planta) at 3rd leaf using 72 h old mycelium of highly virulent isolate (05-06). Control plants were mock inoculated with moist cotton. The experiment was conducted in triplicates. The leaf samples were collected at 24 hpi (hours post inoculation), the necrotic spot was removed and used for protein extraction. The mock inoculated leaves were also collected for the extraction of proteins. Samples from 3 biological replicates were used for the analysis.

### Label free quantitative proteomics

Total leaf protein was extracted (Umadevi and Anandraj 2015) and quantified. Three biological replicates from control and 24 hpi were used to profile the AMPs. For LC-LTQ Orbitrap MS analysis, samples were re-solubilized in 2% [v/v] acetonitrile, 0.1% [v/v] formic acid in water and injected onto an Agilent1200 (Agilent, Santa Clara, CA, USA) nano-flow LC system that was in-line coupled to the nano-electrospray source of a LTQ-Orbitrap discovery hybrid mass spectrometer (Thermo Scientific, San Jose, CA, USA). Peptides were separated on Zorbax 300SB-C18 (Agilent, Santa Clara, CA, USA) by a gradient developed from 2% [v/v] acetonitrile, 0.1% [v/v] formic acid to 80% [v/v] acetonitrile, 0.1% [v/v] formic acid in water over 70 min at a flow rate of 300 nl/min. Full MS in a mass range between m/z 300 and m/z 2000 was performed in an

Orbitrap mass analyzer with a resolution of 30,000 at m/z 400 and an AGC target of  $2 \times 10^5$ . The strongest five signals were selected for CID-MS/MS in the LTQ ion trap at normalized collision energy of 35% using an AGC target of  $1 \times 10^5$  and two micro scans. Dynamic exclusion was enabled with one repeat counts during 45 s and an exclusion period of 120 s. All the 6 samples were included in the analysis where control samples were chosen as reference and all other ion intensity maps from other samples were automatically aligned to the reference. The peptide ion detection method was high resolution. Considering the good initial alignment quality, the data set was not subjected to any further manual correction such as vector editing. Relative quantification using Hi-3 was selected for automatic processing of the software. After successful alignment, no further filtering was applied to subsequent quantification steps in the software. Parameter settings such as no protein grouping and quantitation from non-conflicting features were used for protein building. Peptide identification was performed by CID-based MS/MS of the selected precursors. For protein/peptide identification, MS/MS data were searched against the APD database using an in-house Mascot server (version 2.4) through the ProteomeDiscoverer1.4 software. The search was set up for full tryptic peptides with a maximum of three missed cleavage sites. Carbamidomethyl on cysteine, and oxidized methionine were included as variable modifications. The precursor mass tolerance threshold was 110 ppm, and the maximum fragment mass error was 0.8 Da. The significance threshold of the ion score was calculated based on a false discovery rate of < 1%, estimated by the peptide valid at or node of the Proteome Discoverer software. Ion matching requirements were two fragments per peptide, five fragments per protein, and one peptide per protein. Anova (p)\* 0.05 was kept as significant in selecting the statistically significant fold change expression of AMPs.

### Characterization of AMP sequences

We used APD database for the AMP signature identification (Wang et al. 2016). The AMP signatures were also queried with PhyAMP (Hammami et al. 2009) and CAMPR3 (Waghu et al. 2016). In order to characterize the AMPs in silico, we used the descriptors viz., isoelectric point, aliphatic index and grand average of hydropathy (Gasteiger et al. 2005) (GRAVY) (using ProtParam tool) and the net charge using PhytoAMP database. We also characterized the AMPs as cysteine rich, cysteine free AMPs. The peptide region coding for antigenicity was predicted using Kolaskar and the secondary structures were predicted using GOR4 (Kolaskar and Tongaonkar 1990). Toxin pred (Gupta et al. 2013) and HLP tool (Sharma et al.

2014) was used to predict the toxic peptides and predict the half-life of the peptides.

## Results

### Label free proteomics based identification and expression dynamics of AMPs

The AMP identification was done in uninfected (control) and from the 24 hpi sample. A total of 24 black pepper AMPs (BpAMPs) was matched to the known AMPs from both the samples and the quantitative expression was also deduced using Hi3 method (Table 1). The AMPs mass ranged from 731.4332 to 2340.0385. The relative expression dynamics ranged from 1.3 to 11.15 folds. Fourteen AMPs showed high abundance (5.88–10.59 fold) and the 10 showed low abundance. Low abundance (range) of peptides were found between 3.59 and 1.02 fold. The Hi3 quantification of peptides showed five BpAMPs with above 5 fold increase in expression were BpAMP3 (5.88 fold), BpAMP7 (11.15), BpAMP8 (6.48), BpAMP12 (10.28) and BpAMP23 (10.59).

The identified AMP sequences were queried individually in PhytAMP database using blast tool to identify its plant origin. All the non-plant AMPs were queried against a multi-organism database (CAMP R3) and were found to have matching peptides in this database (Table 2).

### Physiochemical and antimicrobial properties

The number of amino acids and the molecular weight of the BpAMPs ranged from 7–24 to 714–2333.5 respectively. Seven AMPs had an aliphatic index > 70, 10 AMPs < 100 and 6 AMPs < 70 to > 100. GRAVY value is calculated as the sum of hydropathy values of all the amino acids, divided by the number of amino acid residues in the query sequence. Positive and negative GRAVY is an indication of hydrophobicity and hydrophilicity respectively. Among 24 AMPs of black pepper, 12 AMPs were of hydrophobic and another 12 were hydrophilic (Table 3). The net charge of the AMPs varied from 1 to 4. Secondary structure prediction using the GOR 4 secondary structure prediction method showed 23 AMPs having extended strand and random coil in specific proportions. The only AMP (BpAMP17) from black pepper from this study was the type having alpha helix, extended strand and random coil (Table 3). The Kolaskar and Tongaonkar antigenicity prediction was used to determine sequences of antigenic determinants (epitopes) within the AMPs (Table 3). Conserved Domain search for the 24 AMPs showed using NCBI CDD tool identified conserved domains in 2 AMPs. Based on the analysis of cysteine content, the black pepper

AMPs were found to have 13 cysteine free AMPs along with 11 cysteine rich AMPs. The percentage of amino acids in the cysteine free peptides is tabulated (Table 4). The Toxin pred analysis results showed that BpAMP14 was of toxic. Bowman index the protein binding potential was deduced and results in predicting half-life of peptides in the intestine like environment to find the half life for each AMPs and are tabulated (Table 3).

## Discussion

In spite of abundant reports on the isolation of AMPs from many sources, the profile of AMPs expressed/identified from single crop species under certain stress/physiological condition is still unknown. In this study, we aimed to identify AMP using label free proteomic analysis of protein extract from black pepper leaf upon infection with *P. capsici*. The 24 hpi samples were taken from resistant genotype as there is no visible symptom expressed in this cultivar, where as in case of susceptible variety, visible symptoms expresses in 24 hpi (Unpublished data). The peptide data from antimicrobial peptide (APD) database (Wang et al. 2016) was used to query the peptide sequences from control and 24 hpi using Progenesis IQ software. A total of 5 AMPs was found to have similarity to plant peptides and the rest 16 AMPs failed to find a match with the available entries in the PhytAMP database. The search in CAMPR3, the multi-organism database showed the identity for all 16 non-plant AMPs. We suppose them to be homologs of animal /insect AMP signatures. These AMP homologs may be of evolutionary novelty in black pepper.

The differential expression of 24 AMPs suggests that a combinatorial strategy is working in the defense network in black pepper against *P. capsici*. BpAMP3 (AGLQFPVGR) was found to be the Buforin homolog. Buforin isolated from frog showed broad spectrum antimicrobial activity including fungi by penetrating the cell membrane (Park et al. 2000) BpAMP7 (CAPKMKQIGTCGMPQVKCK) was the Hevein type AMP showed similarity to AMP from *Euonymus europaeus* (European spindle tree). This small hevein like chitin binding protein possesses antifungal property and active against *Phytophthora cryptogea*. The chitin-binding hevein-type polypeptides were identified with three (Ac-AMP), four (hevein), five (*Eucommia ulmoides* AMPs) and 10 disulphide bonds (Ee-CBP) from *E. europaeus* (Van den Bergh et al. 2002). The BpAMP7 identified from black pepper was found to have 4 cysteine residues. BpAMP8 (NQCINLEKAR) was identified as Rs-Afp1of Raddish. This type causes membrane permeabilisation and formation of reactive oxygen species (Matejuk et al. 2010) It was demonstrated that that transgenic tomato plants with this AMP was resistant to *Phytophthora*

**Table 1** Quantitative abundance profile of AMPs from black pepper upon 24 hpi of *P. capsici*

Sequence	Peptide ion	Mass	Score	Anova	Fold change	Average normalised abundance (Control)	Average normalised abundance (24 h)	Description	Abundance	Black pepper ID
LGDAVEDLESV GK	4500	1458.7566	56.73	0.30	1.17	2.06e+004	3.17e+004	AP00433	High	BpAMP1
AQSGKTAICKCYVKV CPR	412	2011.0377	36.70	0.54	1.30	2.82e+005	3.44e+005	AP01559	High	BpAMP2
AGLQFPVGR	1113	943.5236	55.33	2.03e-005	5.88	1.24e+004	7.29e+004	AP00307	High	BpAMP3
GICVPIRCPGSMIRQIGTCLGAQVK	2287	2340.0385	20.08	0.36	1.08	1.91e+005	4.40e+004	AP00401	Low	BpAMP4
RGRCLCIGPGVK	1012	1314.7148	21.39	0.92	1.09	6.03e+004	6.55e+004	AP02081	High	BpAMP5
FISGLJGGLMK	1904	1150.6384	31.14	2.47e-005	3.85	2.57e+004	6670.49	AP02274	Low	BpAMP6
CAPKMKQIGTCGMPQVKCKK	287	2226.0018	14.64	6.38e-007	11.15	8861.63	1.64e+005	AP01597	High	BpAMP7
NQCINLEKAR	535	1188.5906	27.55	9.19e-008	6.48	7.38e+004	4.78e+005	AP00286	High	BpAMP8
GAGKAVLGK	14	841.5045	26.89	9.68e-004	3.53	1.21e+006	4.28e+006	AP00383	High	BpAMP9
WLERIGK	191	900.5279	26.31	8.72e-004	2.04	1.65e+005	3.38e+005	AP00777	High	BpAMP10
KNSPFTAKK	1340	1062.5705	25.70	0.35	1.37	2.33e+004	3.20e+004	AP01522	High	BpAMP11
ELENLAAMDLELQK	8301	1616.8157	24.04	3.68e-010	10.28	7467.00	7.67e+004	AP00612	High	BpAMP12
GTCVLVK	4082	760.4121	23.69	1.64e-003	2.53	3594.22	9100.72	AP00891	High	BpAMP13
SCCRSTQARNIYNAPR	4653	1897.8383	22.56	0.01	1.70	2.62e+004	1.54e+004	AP01585	Low	BpAMP14
ALLCKLDK	2792	959.5488	22.39	0.04	1.25	1.66e+004	1.33e+004	AP02299	Low	BpAMP15
TWKRPPQTSCWGIKE	3255	2076.0737	21.99	0.01	3.14	1.72e+004	5481.07	AP01940	Low	BpAMP16
GQVNSACAANCLSLGKAGHCEK	979	2332.0176	21.12	3.10e-004	2.94	1.74e+006	5.93e+005	AP00226	Low	BpAMP17
SSVYGRK	1135	731.4332	21.05	1.83e-003	2.01	2.58e+004	1.28e+004	AP01309	Low	BpAMP18
IQDKEGIPDQQR	4324	1522.7733	20.48	3.30e-006	2.49	3.05e+004	1.23e+004	AP02030	Low	BpAMP19
NKGICVPIR	9157	998.554	17.77	0.85	2.87	1651.76	4735.63	AP00235	Low	BpAMP20
GIKDWIK	4947	900.5037	15.31	4.59e-004	3.59	1.02e+004	2847.57	AP01220	Low	BpAMP21
ALGTLLK	2017	756.4833	12.19	0.16	1.74	8381.04	1.46e+004	AP00909	High	BpAMP22
DAATGLVTGIQS	536	1174.5724	11.16	5.23e-008	10.59	2.40e+004	2.54e+005	AP00699	High	BpAMP23
WVQNYMKHLGRK	4379	1575.7885	10.04	0.06	1.41	3.37e+005	4.74e+005	AP02088	High	BpAMP24

**Table 2** Annotation of black pepper AMPs

AMP	CAMP R3	APD	PhytAMP
BpAMP1	Crystal structure of the hexameric anti-microbial peptide channel dermicidin	Dermicidin	
BpAMP2	Anti microbial peptide ( <i>Aspergillus clavatus</i> ) CAMPSQ2291	AcAMP ( <i>Aspergillus clavatus</i> )	Snakin
BpAMP3	Buforin (CAMPSQ277)	Buforin (Toad)	
BpAMP4	Lingual antimicrobial peptide (defensin family) (SQ1412)	Beta defensin	Ar-AMP Hevein
BpAMP5	NMR structure of CXC chemokine CXCL11/ITAC	Chemokine	GASA-like Snakin
BpAMP6	Maximin-H7 (SQ1780)	Temporin (cationic)	
BpAMP7	Prepro-beta-defensin 1 (SQ2648)	Beta defensin	Ee-CBP leaves (Hevein)
BpAMP8	Gamma-thionin (SQ2567)	Rs-Afp 1 plant defensin	At-AFP1 defensin
BpAMP9	Ponericin-L2 (SQ218)	Ponericin	
BpAMP10	Winter flounder 1 (Pleurocidin family) (CAMPSQ861)	Winter flounder 1	
BpAMP11	Ap (anti fungal) (CAMPSQ3306)	Ap	
BpAMP12	Chrombacin (CAMP SQ2811)	Chrombacin	
BpAMP13	Pilosulin 3 (CAMPSQ495) (from Insect Ant)	Pilosulin 3	
BpAMP14	Pp-AMP1 (defensin) (CAMP SQ3353)	Plant Pp-AMP1 (defensin)	Plant Pp-AMP1 (defensin)
BpAMP15	Brevinin	Brevinin	
BpAMP16	Nigroain-C2 (CAMPSQ3641) from frog	Nigroain C2	
BpAMP17	Defensin-1 ( <i>Apis mellifera carnica</i> ) (CAMPSQ4363)	Royalisin	
BpAMP18	No hit	Odorranain	
BpAMP19	CgUbiquitin (CAMPSQ3702)	Cg ubiquitin	
BpAMP20	LAP-like antimicrobial peptide (fragment) (defensin) (CAMPSQ 6679)	Beta defensin	
BpAMP21	Ascaphin-5(human erythrocytes) (CAMPSQ4333)	Ascaphin 5	
BpAMP22	Dermatoxin S1 (frog) (CAMPSQ 2946)	Dermatoxin	
BpAMP23	Dahlein 4.3 (synthetic construct) (CAMPSQ2851)	Dahlein	
BpAMP24	CCL 13	CCL 13 chemokine	

*infestans* (Parashina et al. 2000). BpAMP12 (ELEN-LAAMDLELQK) was the chrombacin analog. These peptides have the ability to induce chemotaxis and initiation of release of cytokines (Salzet and Stefano 2003). BpAMP23 (DAATGLVTGIQS) was found to be an analog of Dahlein, bioactive peptides from frog *Litoria dahlia* which was found to have wide spectrum activity (Wegener et al. 2001).

Isoelectric point is an important factor as it affects the solubility of the AMP. The black pepper AMPs had pI range from 3.8 to 10.3 denoting the presence of most acidic and alkaline range having peptides. Aliphatic index shows the thermal stability of the AMPs. The aliphatic index of 16 black pepper AMPs were with 70–100 showing greater thermal stability.

Studies demonstrated that at least net charge 2 is required for the amphipathic nature (Hancock 1997). The net charge of most of the black pepper AMPs was found to be 2 and above, which indicated their antimicrobial potential. The total of ionizable amino acid residues at a

particular pH determines the anionic or cationic net surface charge to AMPs. Anionic /cationic AMPs are constitutive or inducible defense barriers against microbial infections and also they might have the ability to improve host immunity by acting as immune modulators (Robinson et al. 2012). We found 4 anionic AMPs (BpAMP1, 12, 19 and 23) while another 18 AMPs were cationic in nature. The plant derived anionic AMPs are attractive molecule against cancer. This group of AMPs is reported as host defense peptides from plants and is shown to have anticancer property (Song et al. 2012). On the other hand, the plant cationic AMPs are shown to have activity against negatively charged microbial membranes.

The majority of black pepper AMPs were found to have extended strand and random coil secondary structures. The reports state that the anionic peptides should have extended strand and random coil (Powers and Hancock 2003). The extended class of peptides is rich in proline and/or glycine contents and lacks classical secondary structures. The random coils are found to be involved in cell permeation in

**Table 3** Physicochemical/antimicrobial properties of black pepper AMPs

AMP	Number of amino acids	MW	pI	Net charge	Aliphatic index	Secondary structure (ES: extended strand; RC: random coil; AH: alpha helix)	Bowman index(kcal/mol)	Intestinal half life(sec)	Toxicity	Most antigenic region	(GRAVY)
BpAMP1	14	1459.6	4.32	-2	104.29	ES (50%); RC (50%)	1.71	0.766	Non-toxin	VEDLESV	-0.4
BpAMP2	18	1955.3	9.42	4	65	Extended strand (38.89%); random coil (61.11)	1.21	1.620	Non-toxin	CYVKVCP	-0.028
BpAMP3	9	944.1	9.79	1	86.67	Extended strand (44.44%); random coil (55.56%)	0.53	1.328	non-toxin	AGLQFPV	0.244
BpAMP4	24	2488	9.22	3	93.33	Extended strand (41.67%); random coil (58.33%)	0.53	2.710	Non-toxin	ICVPIRC	0.446
BpAMP5	12	1258.5	9.7	3	89.17	Extended strand (58.33%); random coil (41.67%)	1.34	2.172	Non-toxin	CLCIGPG	0.15
BpAMP6	11	1135.4	8.75	1	141.82	Extended strand (54.55%); random coil (45.45%)	-1.71	1.675	Non-toxin	FISGLIG	1.4
BpAMP7	20	2154.7	9.21	4	39	Extended strand (45%); random coil (55%)	0.66	2.385	Non-toxin	CGMPQVK	-0.15
BpAMP8	10	1188.3	8.22	1	88	Extended strand (20%); random coil (80%)	3.31	0.617	Non-toxin	CINLEKA	-0.98
BpAMP9	9	799.9	10	2	97.78	Extended strand (55.56%); random coil (44.44%)	-0.47	0.859	Non-toxin	AGKAVLG	0.289
BpAMP10	7	901	8.75	1	111.43	Extended strand (71.43%); random coil (28.57%)	2.02	1.129	Non-toxin	WLERIGK	-0.7
BpAMP11	9	1020.2	10.3	3	111.11	Extended strand (44.44%); random coil (55.56%)	2.71	1.111	Non-toxin	SPFTAKK	-1.522
BpAMP12	14	1616.8	4	-3	125.71	Extended strand (21.43%); random coil (42.86%)	1.51	0.106	Non-toxin	AAMDLEL	-0.3
BpAMP13	7	718.9	8.22	1	138.57	Extended strand (71.43%); random coil (28.57%)	-1.01	0.746	Non-toxin	GTCVLVK	1.386
BpAMP14	16	1840	9.69	3	36.88	Extended strand (31.25%); random coil (68.75%)	3.87	0.427	Toxin	CCRSTQA	-1.006
BpAMP15	8	903.1	8.24	1	158.75	Extended strand (25%); random coil (75%)	0.24	2.604	Non-toxin	ALLCKLD	0.55
BpAMP16	17	2077.4	9.3	2	45.88	Extended strand (23.53%); random coil (76.47%)	1.6	1.834	Non-toxin	PPQTSC	-0.741
BpAMP17	24	2333.5	6.72	1	61.25	Alpha helix (25%); Extended strand (8.33%); random coil (75%)	1.17	2.740	Non-toxin	ACAANCL	-0.217
BpAMP18	7	731.8	11	2	82.86	Extended strand (57.14%); random coil (42.86%)	2.02	0.594	Non-toxin	SSVVGRK	-0.286
BpAMP19	13	1523.6	4.56	-1	60	Extended strand (15.38%); random coil (84.62%)	3.88	1.097	Non-toxin	GIPPDQQ	-1.846
BpAMP20	9	999.2	9.51	2	118.89	Extended strand (44.44%); random coil (55.56%)	1.22	1.107	Non-toxin	KGICVPI	0.2
BpAMP21	7	859	8.59	1	111.43	Extended strand (28.57%); random coil (71.43%)	0.95	1.143	Non-toxin	GIKDWIK	-0.514
BpAMP22	7	714.9	8.8	1	181.43	Extended strand (57.14%); random coil (42.86%)	-1.34	1.738	Non-toxin	ALGTLLK	1.171
BpAMP23	12	1132.2	3.8	-1	105.83	Extended strand (58.33%); random coil (41.67%)	0.28	0.543	Non-toxin	LVTGIQS	0.508
BpAMP24	12	1559.8	10.29	4	56.67	Extended strand (25%); random coil (75%)	2.36	1.612	Non-toxin	VQNYMKH	-1.267




**Table 4** Cysteine free AMPs with the amino acid composition and regulation

AMP	Amino acid composition (%)	Regulation upon <i>Phytophthora</i> 24h (fold change)
BPAMP1	Ala, Ser (7.14); ASP, Glu, Gly, Val, Lys, Leu (14.29)	Up (1.17)
BPAMP3	Ala, Phe, Pro, Gln, Arg, Val (11.11); <b>Gly (22.22)</b>	Up (5.88)
BPAMP6	Phe, Lys, Met, Ser (9.09);Ile, Leu (18.18); <b>Gly (27.27)</b>	Down (3.85)
BPAMP9	Ala, Lys (22.22); <b>Gly (33.33)</b> ; Leu, Val (11.11)	Up (3.53)
BPAMP10	Glu, Gly, Ile, Leu, Lys, Arg, Try (14.29)	Up (2.04)
BPAMP11	Ala, Phe, Asn, Pro, Ser, Thr (11.11); <b>Lys (33.33)</b>	Up (10.28)
BPAMP12	Ala (14.29); Asp, Lys, Met, Asn, Gln (7.14); <b>Glu (21.43)</b> ; Leu ( <b>28.47</b> )	Up (2.53)
BPAMP18	Gly, Lys, Arg (14.29); <b>Ser, Val (28.57)</b>	Down (2.01)
BPAMP19	Asp, Ile, pro (15.38); <b>Gln(23.08)</b> ; Glu, Gly, Lys, Arg (7.69)	Down (2.49)
BPAMP21	Asp, Gly, Trp (14.29); <b>Ile, Lys (28.57)</b>	Down (3.59)
BPAMP22	Ala, Gly, Lys, Thr (14.29); <b>Leu (42.86)</b>	Up (1.74)
BPAMP23	<b>Ala, Gly, Thr (16.67)</b> ; Asp, Ile, Leu, Gln, Ser, Val (8.33)	Up (10.59)
BPAMP24	Gly, His, Leu, Met, Asn, Gln, Arg, Val, Trp, Tyr (8.33); <b>Lys (16.67)</b>	Up (1.41)

case of Indolicidin AMP (Zhang et al. 2001). The conserved domain search yielded BpAMP4 (GICVPIRCPGSMRQIGTCLGAQVK) with defensin beta superfamily and BpAMP (IQDKEGIPPDQQR) with UBQ super family conserved domain (Fig. 1). These results further strengthen that the label free proteomics approach as reliable and quick method to identify AMPs even from complex samples and possibility to find the gene fragment coding for AMPs.

The cysteine free AMPs are more common in animal, insects and they were found to be linear peptides without cysteine with a high proportion of certain residues. In plant, till now only 3 reports are available on linear cysteine free AMPs (Egorov et al. 2005; Zipfel 2009; Silva et al. 2012). Out of 13 cysteine free BpAMPs, 9 AMPs were found to be in up-regulation (BpAMP1, 3, 9, 10, 11, 12, 22, 23 and 24) and the remaining 4 were with down-regulation. The cysteine rich AMPs are very common to plant kingdom with



**Fig. 1** Black pepper AMPs with conserved domains

varying number of cysteine residues. In this study, we identified AMPs in black pepper with 4 cysteine residues (BpAMP7), 3 cysteine residues (BpAMP2, 4 and 17), 2 cysteine residues (BpAMP5 and 14) and 1 cysteine residue (BpAMP8, 13, 15, 16 and 20). Among the 11 cysteine rich AMPs, two were up-regulated upon *Phytophthora* infection, BpAMP7 (11.15 fold) that is similar to hevein type AMP and BpAMP8 (6.48 fold) similar to Rs-Afp-1, suggesting them to be the effective candidate AMPs as molecule against *Phytophthora*.

In addition to this, the analysis to detect the toxic nature and half-life of the AMPs which are important for any drug development. This information would be important for any researcher to use the peptides towards drug development.

By using label free proteomics strategy, we established for the first time the black pepper peptidome associated with the innate immunity against *Phytophthora*. We showed the occurrence of both cysteine rich, cysteine free AMPs from a complex sample and some major AMP signatures as innate immunity factors against *Phytophthora*. However, whether all the AMPs or some major AMPs are contributing to the pathogen resistance in this genotype still needs to be worked out. Our work presented here will offer a basic platform for further studying the immunology and evolutionary significance of these newly discovered AMPs in black pepper and also utilizing some of the AMPs as next generation fungicide molecules.

**Acknowledgements** We thank the Indian Council of Agricultural Research, New Delhi for funding through Outreach program on *Phytophthora*, *Fusarium* and *Ralstonia* diseases of horticultural and field crops (PhytoFuRa) and mass spectrometry facility, C-CAMP, NCBS, Bangalore for the LC/MS analysis.

#### Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

## References

Anandaraj M (2000) Diseases of black pepper. In: Ravindran PN (ed) Black pepper (*Piper nigrum* L.). Harwood Academic Publishers, New York, pp 239–268

Asiegbu FO, Choi W, Li G, Nahalkova J, Dean RA (2003) Isolation of a novel antimicrobial peptide gene (SpAMP) homologue from *Pinus sylvestris* (Scots pine) following infection with the root rot fungus *Heterobasidion annosum*. FEMS Microbiol Lett 228:27–31

Cammue BPA, De Bolle MFC, Terras FRG, Proost P, Van Damme J, Rees SB, Vanderleyden J, Broekaert WF (1992) Isolation and characterization of a novel class of plant antimicrobial peptides from *Mirabilis jalapa* L. seeds. J Biol Chem 267:2228–2233

Egorov TA, Odintsova TI, Vitaliy A, Pukhalsky VA, Grishin EV (2005) Diversity of wheat anti-microbial peptides. Peptides 26:2064–2073

Fan B, Shen L, Liu K, Zhao D, Yu M, Sheng J (2008) Interaction between nitric oxide and hydrogen peroxide in post harvest tomato resistance response to *Rhizopus nigricans*. J Sci Food Agric 88:1238–1244

Gasteiger E, Hoogland C, Gattiker A (2005) Protein identification and analysis tools on the ExPASy server. In: Walker JM (ed) The proteomics protocols handbook. Humana Press, New York, pp 571–607

Gupta S, Kapoor P, Chaudhary K, Gautam A, Kumar R, Raghava GPS (2013) In silico approach for predicting toxicity of peptides and proteins. PLoS ONE 8(9):e73957

Hammami R, Ben Hamida J, Fliss I (2009) PhytAMP: a database dedicated to antimicrobial plant peptides. Nucleic Acids Res D 963:8

Hancock RE (1997) Peptide antibiotics. Lancet 349(9049):418–422

Ke T, Cao H, Huang J, Hu F, Huang J, Dong C, Ma X, Yu J, Mao H, Wang X, Niu Q, Hui F, Liu S (2015) EST-based in silico identification and in vitro test of antimicrobial peptides in *Brassica napus*. BMC Genom 16:653

Kolaskar AS, Tongaonkar PC (1990) A semi-empirical method for prediction of antigenic determinants on protein antigens. FEBS Lett 276(1–2):172–174

Matejuk A, Leng Q, Begum MD, Woodle MC, Scaria P, Chou S-T, Mixson AJ (2010) Peptide-based antifungal therapies against emerging infections. Drugs Future 35(3):197

Nawrot R, Baryliski J, Nowicki G, Broniarczyk J, Buchwald W, Gozdzicka-Jozefiak A (2014) Plant antimicrobial peptides. Folia Microbiol 59(3):181–196

Parashina EV, Serdobinskii LA, Kalle EG, LAVOROVA NV, AVETNOV VA, LUMIN VG, NARODITSKII BS (2000) Genetic engineering of oilseed rape and tomato plants expressing a radish defensin gene. Russ J Plant Physiol 47:417–423

Park CJ, Park CB, Hong SS, Lee HS, Lee SY, Kim SC (2000) Characterization and cDNA cloning of two glycine- and histidine-rich antimicrobial peptides from the roots of shepherd-spurge, *Capsella bursa-pastoris*. Plant Mol Biol 44:187–197

Pelegri PB, Del Sarito RP, Silva ON, Franco OL, Grossi-de-sa MF (2011) Antimicrobial peptides from plants: what they are and how they probably work. Biochem Res Int 2011:250349

Powers JP, Hancock RE (2003) The relationship between peptide structure and antibacterial activity. Peptides 24(11):1681–1691

Rahnamaeian M (2011) Antimicrobial peptides: modes of mechanism, modulation of defense responses. Plant Signal Behav 6:1325–1332

Robinson MW, Hutchinson AT, Donnelly S (2012) Antimicrobial peptides: utility players in innate immunity. Front Immunol 3:325

Salzet M, Stefano G (2003) Chromacin-like peptide in leeches. Neuro Endocrinol Lett 24(3/4):227–232

Sarika, Iqebalb MA, Rai A (2012) Biotic stress resistance in agriculture through antimicrobial peptides. Peptides 36:322–330

Scott MG, Dullaghan E, Mookherjee N, Waldbrook M, Thompson A, Wang A, Lee K, Doria S, Hamil P, Yu JJ, Li Y, Donini O, Guarna MM, Finlay BB, North JR, Hancock RE (2007) An anti infective peptide that selectively modulates the innate immune response. Nat Biotechnol 25:465–472

Sharma A, Singla D, Rashid M, Raghava GPS (2014) Designing of peptides with desired half-life in intestine-like environment. BMC Bioinformatics 15:282

Silva ON, Porto WF, Migliolo L, Mandal SM, Gomes DG, Holanda HH, Silva RS, Dias SC, Costa MP, Costa CR, Silva MR, Rezende TM, Franco OL (2012) Cn-AMP1: a new promiscuous peptide with potential for microbial infections treatment. Biopolymers 98(4):322–331



- Song R, Wei R, Luo H, Wang D (2012) Isolation and characterization of an antibacterial peptide fraction from the pepsin hydrolysate of half-fin anchovy (*Setipinna taty*). *Molecules* 17:2980–2991
- Umadevi P, Anandaraj M (2015) An efficient protein extraction method for proteomic analysis of black pepper (*Piper nigrum* L.) and generation of protein map using nano LC-LTQ Orbitrap mass spectrometry. *Plant Omics* 8(6):500–507
- Van den Bergh KPB, Proost P, Van Damme J, Coosemans J, Van Damme EJM, Peumans WJ (2002) Five disulfide bridges stabilize a hevein-type antimicrobial peptide from the bark of spindle tree (*Euonymus europaeus* L.). *FEBS Lett* 530(1–3):181–185
- Waghu FH, Barai RS, Gurung P, Idicula-Thomas S (2016) CAMPR3: a database on sequences, structures and signatures of antimicrobial peptides. *Nucleic Acids Res* 44:D1094–D1097
- Wang G, Li X, Wang Z (2016) APD3: the antimicrobial peptide database as a tool for research and education. *Nucleic Acids Res* 44:D1087–D1093
- Wegener KL, Brinkworth CS, Bowie JH (2001) Bioactive dahlein peptides from the skin secretions of the Australian aquatic frog *Litoria dahlii*: sequence determination by electrospray mass spectrometry. *Rapid Commun Mass Spectrom* 15(18):1726–1734
- Weinhold A, Wielsch N, Svatos A, Baldwin IT (2015) Label-free nanoUPLC-MS<sup>E</sup> based quantification of antimicrobial peptides from the leaf apoplast of *Nicotiana attenuate*. *BMC Plant Biol* 15:18
- Zhang L, Rozek A, Hancock REW (2001) Interaction of cationic antimicrobial peptides with model membranes. *J Biol Chem* 276(38):35714–35722
- Zhou M, Hu Q, Li Z, Li D, Chen C, Luo H (2011) Expression of a novel antimicrobial peptide Penaeidin 4-1 in creeping bentgrass (*Agrostis stolonifera* L.) enhances plant fungal disease resistance. *PLoS ONE* 6(9):e24677
- Zipfel C (2009) Early molecular events in PAMP-triggered immunity. *Curr Opin Plant Biol* 12:414–420