

Genetic characterization and fine mapping of the novel *Phytophthora* resistance gene in a Chinese soybean cultivar

Jiqing Zhang · Changjian Xia · Xiaoming Wang ·
Canxing Duan · Suli Sun · Xiaofei Wu ·
Zhendong Zhu

Received: 5 July 2012 / Accepted: 23 February 2013 / Published online: 7 March 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract *Phytophthora* root rot (PRR), caused by *Phytophthora sojae* Kaufmann & Gerdemann, is one of the most destructive diseases of soybean [*Glycine max* (L.) Merr.]. Deployment of resistance genes is the most economical and effective way of controlling the disease. The soybean cultivar ‘Yudou 29’ is resistant to many *P. sojae* isolates in China. The genetic basis of the resistance in ‘Yudou 29’ was elucidated through an inheritance study and molecular mapping. In response to 25 *P. sojae* isolates, ‘Yudou 29’ displayed a new resistance reaction pattern distinct from those of differentials carrying known *Rps* genes. A population of 214 F_{2:3} families from a cross between ‘Jikedou 2’ (PRR susceptible) and ‘Yudou 29’ was used for *Rps* gene mapping. The segregation fit a ratio of 1:2:1 for resistance:segregation:susceptibility within this population, indicating that resistance in ‘Yudou 29’ is controlled by a single dominant gene. This gene was temporarily named *RpsYD29* and mapped on soybean chromosome 03 (molecular linkage group N; MLG N) flanked by SSR markers SattWM82-50 and Satt1k4b at a genetic distance of 0.5 and 0.2 cM, respectively. Two nucleotide binding site-leucine rich repeat (NBS-LRR)

type genes were detected in the 204.8 kb region between SattWM82-50 and Satt1k4b. These two genes showed high similarity to *Rps1k* in amino acid sequence and could be candidate genes for PRR resistance. Based on the phenotype reactions and the physical position on soybean chromosome 03, *RpsYD29* might be a novel allele at, or a novel gene tightly linked to, the *Rps1* locus.

Introduction

Phytophthora root rot (PRR), caused by *Phytophthora sojae* Kaufmann & Gerdemann, is one of the most destructive diseases of soybean [*Glycine max* (L.) Merr.] (Bernard et al. 1957; Kuan and Erwin 1980). This disease has been reported in most soybean-production regions of the world, since it was first noted in America in 1948 (Tyler 2007; Wrather et al. 2001). In China, PRR was first observed in 1991 in Heilongjiang Province (Shen and Su 1991). Since then, the disease has become widespread in Heilongjiang Province, and the annual infected area is estimated at over 150,000 ha (Zhang et al. 2010). It has now also spread to Fujian Province (Chen et al. 2004).

PRR can be effectively controlled through the use of complete or partial resistance genes (Sugimoto et al. 2012). Partial resistance to *P. sojae* can be ineffective when plants are subjected to high disease pressure (Dorrance et al. 2003), so the deployment of race-specific, complete resistance genes in soybean cultivars has been the primary method used to control PRR (Schmitthenner 1999). To date, 14 genes conferring resistance to *P. sojae* (*Rps*) have been identified at 8 loci, *Rps1* to *Rps8*, which were mapped to molecular linkage groups (MLG) N, J, F, G, G, G, N and F, respectively (Sandhu et al. 2005). Five functional genes *Rps1a*, *Rps1b*, *Rps1c*, *Rps1d* and *Rps1k* have been mapped at the *Rps1*

Communicated by D. Lightfoot.

Electronic supplementary material The online version of this article (doi:10.1007/s00122-013-2073-1) contains supplementary material, which is available to authorized users.

J. Zhang · C. Xia · X. Wang · C. Duan · S. Sun · X. Wu ·
Z. Zhu (✉)

MOA Key Lab of Soybean Biology (Beijing), the National Key Facility for Crop Gene Resources and Genetic Improvement, Institute of Crop Science, Chinese Academy of Agricultural Sciences, 12 Zhongguancun South Street, Beijing 100081, People’s Republic of China
e-mail: zhuzhendong@caas.cn

locus (Bernard et al. 1957; Buzzell and Anderson 1992; Mueller et al. 1978), and three alleles *Rps3a*, *Rps3b* and *Rps3c* have been mapped at the *Rps3* locus (Mueller et al. 1978; Ploper et al. 1985). Additionally, five *Rps* genes, *RpsYB30*, *RpsYD25* (*Rpsyu25*), *RpsZS18*, *RpsSN10* and the *Rps* gene in soybean cultivar ‘Waseshiroge’, were mapped to MLG L, N, D1b, F and N, respectively (Fan et al. 2009; Sugimoto et al. 2011; Sun et al. 2011; Yao et al. 2010; Yu et al. 2010; Zhu et al. 2007).

Among these *Rps* genes, *Rps1k* is of great interest because it confers stable, broad-spectrum resistance in the USA (Schmitthenner et al. 1994), although new *Rps1k*-virulent *P. sojae* isolates have been reported in some areas (Dorrance et al. 2003; Schmitthenner 1999). This gene was first identified in cultivar ‘Kingwa’ (Bernard and Cremeens 1981) and has been transferred to soybean cultivars in the USA (Slaminko et al. 2010). The BAC-cloned *Rps1k* region contains two classes of functional coiled coil-nucleotide binding-leucine rich repeat genes (*Rps1k-1* and *Rps1k-2*) and repetitive sequences (Gao et al. 2005; Gao and Bhattacharyya 2008). In addition to *Rps1k*, the *Rps1c* and *Rps1a* genes are also frequently introduced into commercial cultivars in the USA (Slaminko et al. 2010). However, the continuous use of the *Rps* genes in soybean cultivars has created selection pressure for the evolution of new pathogenic *P. sojae* isolates that can overcome the resistance conferred by these genes.

Since 1955, at least 55 *P. sojae* races with virulence pathotype varying from 7 to 1a, 1b, 1c, 1d, 1k, 3a, 6 and 7 were identified in the USA (Grau et al. 2004). *P. sojae* also presents similarly abundant virulence diversity in China (Zhang et al. 2010; Zhu et al. 2003). So far, 12 *P. sojae* races (1, 3, 4, 5, 9, 11, 13, 17, 21, 24, 44 and 54) were found in Heilongjiang Province, China (Zhang et al. 2010). In addition, other virulence pathotypes that differ from the 55 *P. sojae* races in the USA were also found in China (Zhu et al. 2003). Most of the soybean cultivars or germplasm containing the known *Rps* genes, except *Rps1a*, *Rps1c* and *Rps1k*, are not effective at controlling PRR in the major soybean-production regions (Cui et al. 2010; Tang et al. 2010; Zhang et al. 2010; Zhu et al. 2003). Although new races of *P. sojae* appear with the release of resistant cultivars, the use of genetic resistance still remains the most effective strategy to reduce losses caused by the pathogen. Thus, it is necessary to identify new *Rps* genes and develop molecular markers to genes of interest for marker-assisted selection (MAS).

In our previous study, we found that the cultivar ‘Yudou 29’ had a broad-spectrum resistance (Chen et al. 2008). The objectives of this study were to further characterize the inheritance of the *Rps* gene(s) in the cultivar ‘Yudou 29’, identify SSR markers for fine mapping and predict the candidate gene(s).

Materials and methods

Plant materials

‘Yudou 29’ is a PRR-resistant cultivar, and ‘Jikedou 2’ is a PRR susceptible cultivar. A population of 214 families derived from a cross between ‘Yudou 29’ and ‘Jikedou 2’ was used for gene mapping. The F₁ plants from the cross were self-pollinated to produce a population of F₂ plants. Each F₂ plant was self-pollinated and threshed individually to yield seeds of the F_{2:3} families for both genotype and phenotype evaluations.

To determine which *Rps* gene or *Rps* gene combination was present in ‘Yudou 29’, phenotype analysis was performed using 15 differentials carrying a single *Rps* gene: Harlon (*Rps1a*), Harosoy13XX (*Rps1b*), Williams79 (*Rps1c*), PI103091 (*Rps1d*), Williams82 (*Rps1k*), L76-988 (*Rps2*), L83-570 (*Rps3a*), PRX146-36 (*Rps3b*), PRX145-48 (*Rps3c*), L85-2352 (*Rps4*), L85-3059 (*Rps5*), Harosoy62XX (*Rps6*), Harosoy (*Rps7*), PI399073 (*Rps8*) and Yudou25 (*RpsYD25*). Williams (*rps*) was used as the susceptible check to indicate successful inoculation.

Phytophthora sojae inoculation and PRR evaluation

A total of 25 *P. sojae* isolates with different virulence were used in the phenotype test (Table 1). For each isolate inoculation, 12 seeds of each differential were planted per replication in a pot with a diameter of 10 cm, and all differentials were tested in 3 replications using the hypocotyl-inoculation technique (Haas and Buzzell 1976). To test the phenotypes of the population, the isolate PsMC1 was used to inoculate ‘Yudou 29’, ‘Jikedou 2’, and 30 individual seedlings of each F_{2:3} family using the same technique as above.

After inoculation, the plants were placed in a mist room with relative humidity 100 % and an average temperature of 25 °C for 2 days. They were then placed in a greenhouse with an average temperature of 25 °C. Reactions were recorded as the percentage of dead seedlings in each F_{2:3} family at 6 days post-inoculation (DPI). In accordance with Gordon et al. (2006), families with 0–20 % dead seedlings were scored as homozygous resistant (R), families with 80–100 % dead seedlings were considered homozygous susceptible (S), and families with 21–79 % of dead seedlings were scored as segregating (Rs).

DNA preparation and pooling for bulk segregation analysis

DNA was extracted from soybean leaf tissues collected from 15–30 seedlings of each family using the CTAB method with minor modifications (Allen et al. 2006). Resistant and susceptible bulks for the bulk segregation analysis (BSA)

Table 1 Virulence pathotype of 25 *P. sojae* isolates from China classified using 15 differentials carrying the *Rps1a*, *Rps1b*, *Rps1c*, *Rps1d*, *Rps1k*, *Rps2*, *Rps3a*, *Rps3b*, *Rps3c*, *Rps4*, *Rps5*, *Rps6*, *Rps7*, *Rps8* and *RpsYD25* gene, respectively

| Isolate | Virulence pathotype | Isolate | Virulence pathotype |
|---------|-------------------------------|----------|--|
| PsJL1-1 | 1a, 1c | PsTA3 | 1b, 3a, 3c, 4, 5, 6, 7, 8 |
| PsJMS3 | 1a, 2 | PsJS9 | 1d, 3a, 3b, 3c, 4, 5, 6, 8 |
| PsHJL1 | 3c, 7, YD25 | Ps41-1 | 1a, 1d, 2, 3b, 3c, 5, 7, 8 |
| PsAH4 | 2, 3a, 4, 5 | PsJS7 | 1a, 1d, 3a, 3b, 3c, 4, 5, 6 |
| PsJL4-1 | 1a, 1b, 1c, 1k | PsJL5 | 1a, 1b, 1c, 1d, 2, 3b, 3c, 7, 8 |
| PsJL3-2 | 1a, 1b, 1c, 1k, 7 | PsJS8 | 1b, 1d, 3a, 3b, 3c, 4, 5, 6, 7, 8 |
| PsFJ | 1b, 4, 5, 6, 7, YD25 | Ps52 | 1b, 1d, 3c, 4, 5, 6, 7, 8, YD25 |
| PsSX1 | 1a, 1b, 1c, 1d, 1k, 7 | Ps53 | 1a, 1b, 1d, 2, 3b, 3c, 5, 7, YD25 |
| PsAH3 | 1a, 1k, 3a, 4, 5, 6, 7 | PsMC1 | 1a, 1c, 1k, 2, 3b, 3c, 4, 5, 6, 7, 8 |
| PsXJ | 1a, 1d, 2, 3b, 3c, 7, 8 | PsJN4 | 1b, 1c, 1d, 2, 3b, 3c, 4, 5, 6, 7, 8 |
| PsAH1 | 1b, 1c, 2, 3a, 4, 5, 6, 8 | PsAH5 | 1a, 1b, 1d, 2, 3a, 3c, 4, 5, 6, 7, 8, YD25 |
| PsBr1 | 1a, 1b, 4, 5, 6, 8, YD25 | PsHY33-1 | 1a, 1k, 2, 3a, 3b, 3c, 4, 5, 6, 7, 8, YD25 |
| PsAM1 | 1b, 1c, 1d, 1k, 2, 6, 7, YD25 | | |

were prepared from DNA samples of either ten homozygous-resistant or ten homozygous-susceptible $F_{2:3}$ families (Michelmore et al. 1991). DNA bulks were prepared by pooling 1 μ g DNA of each selected family. The final concentration of the DNA bulks was adjusted to 50 ng/ μ l.

SSR marker development and analysis

The *Rps* gene in ‘Yudou 29’ was first mapped with the SSR markers described in Soybase (<http://soybase.org>). Since the *G. max* sequences (<http://www.phytozome.net/soybean>) have been released (Schmutz et al. 2010), new markers could be developed to finely map the gene. Thus, new SSR markers between Sat_186 and Satt530 were designed with Primer Premier 5.0 (Premier Biosoft International, Palo Alto, CA, USA) with default parameters according to the sequence downloaded from Phytozome (<http://www.phytozome.net/soybean>), in which the simple repeat sequences were identified by SSR Hunter 1.3 (<http://en.bio-soft.net/dna/SSRHunter>). Because *Rps1k* is located at the *Rps1* locus on MLG N (chromosome 03) and has been sequenced, new SSR markers were also designed from the contig containing the *Rps1k-1*, *Rps1k-2* and *Rps1k-3* (GenBank accession no. EU450800).

PCR reactions were performed on a thermal cycler (Biometra, USA) in 20 μ l reaction volumes containing 50 ng genomic DNA, 2.5 μ l 10 \times PCR reaction buffer (2.0 mM MgCl₂), 0.2 mM each dNTP, 1.0 U of Taq DNA polymerase and 0.2 μ M of each primer. The PCR amplification consisted of an initial denaturation at 94 $^{\circ}$ C for 5 min, followed by 35 cycles of denaturation at 94 $^{\circ}$ C for 45 s, annealing at 49–52 $^{\circ}$ C for 45 s and extension at 72 $^{\circ}$ C for 45 s, and final extension at 72 $^{\circ}$ C for 10 min. PCR products were mixed with 4 μ l of 6 \times loading buffer (0.25 % bromophenol blue, 0.25 % xylene cyanol FF and 40 % sugar) and were separated by 6 % polyacrylamide

sequencing gel. The BSA polymorphic SSR markers were further tested for the entire $F_{2:3}$ mapping population.

Data analysis and linkage map construction

Resistant (R) and intermediate (I) phenotypes to the isolates (resistant reactions) were recorded as “1” for presence, and susceptible phenotypes were recorded as “0”. Cluster analysis was performed with different modules of the NTSYS-PC software, version 2.11L (Rohlf 2000).


The segregation patterns of phenotypes and selected SSR markers in the mapping population were tested for goodness-of-fit to Mendelian segregation ratio with Chi-square (χ^2) analysis. A genetic linkage map of *RpsYD29* was constructed with the Joinmap 4.0 linkage analysis software (Van Ooijen 2006). Linkage groups were determined using a log-likelihood (LOD) threshold of 3.0.

Results

Phenotype reaction of ‘Yudou 29’ to *P. sojae* isolates

‘Yudou 29’ was resistant to 19 of the 25 isolates tested, while the 15 differentials carrying a single known *Rps* gene were resistant to 8–18 isolates. The reactions of ‘Yudou 29’ differed from those of Harlon, Harosoy13XX, Williams79, PI103091 and Williams82 to 17, 10, 10, 11 and 13 isolates, respectively. The reaction pattern of ‘Yudou 29’ to the 25 *P. sojae* isolates did not resemble any of the 15 differentials. The genetic diversity and phenotypic relationships between ‘Yudou 29’ and the 15 differentials in this study were depicted by cluster analysis (Fig. 1). Cutting at the coefficient 0.726, ‘Yudou 29’ comprised a single subgroup, which suggested that it might contain a novel *Rps* gene.

Fig. 1 Dendrogram revealed by UPGMA cluster analysis of the Phytophthora resistance among ‘Yudou 29’, 15 differentials carrying a single *Rps* gene: Harlon (*Rps/a*), Harosoy13XX (*Rps/b*), Williams79 (*Rps/c*), PI103091 (*Rps/d*), Williams82 (*Rps/k*), L76-988 (*Rps/2*), L83-570 (*Rps/3a*), PRX146-36 (*Rps/3b*), PRX145-48 (*Rps/3c*), L85-2352 (*Rps/4*), L85-3059 (*Rps/5*), Harosoy62XX (*Rps/6*), Harosoy (*Rps/7*), PI399073 (*Rps/8*) and Yudou25 (*RpsYD25*) and the susceptible cultivar Williams



Phenotype analysis for the mapping population

‘Jikedou 2’ plants inoculated with PsMC1 displayed severe rot at the inoculation site and all plants were dead at 6 DPI. ‘Yudou 29’ plants showed no symptoms at 6 DPI, and thus were resistant to the isolate PsMC1.

Among the 214 $F_{2:3}$ families of the mapping population, 51 were homozygous resistant (R), 60 were homozygous susceptible (S) and 103 were segregating (Rs) to the isolate PsMC1 (Table 2). A segregation ratio of 51:103:60 in the $F_{2:3}$ population fits well with the genetic model ratio 1:2:1 ($\chi^2 = 1.06$, $p = 0.62$). This result suggested that Phytophthora resistance in ‘Yudou 29’ was controlled by a single dominant gene, which was temporarily named *RpsYD29*.

Molecular mapping of the *RpsYD29* gene

Using the BSA method, five SSR markers, Satt152, Satt159, Sat_186, Satt631 and Satt530 on chromosome 03 (MLG N), showed polymorphisms between ‘Yudou 29’

and ‘Jikedou 2’, as well as between the resistant bulk and the susceptible bulk. Linkage analysis further revealed that *RpsYD29* was linked to these five SSR markers and located between Sat_186 and Satt530 (Fig. 2).

A 2.34 Mb DNA segment of soybean chromosome 03 between Sat_186 and Satt530 was identified using a “BLAST genome” search in the Phytozome soybean genome database. Within this region, 218 SSR loci were identified by SSR hunter. A total of 156 SSR loci with 15 bp or greater repeat motifs were selected to design primers for fine mapping of *RpsYD29*. Three SSR markers, SattWM82-32, SattWM82-39 and SattWM82-50, showed polymorphisms between ‘Yudou 29’ and ‘Jikedou 2’ (Supplementary Table 1).


In addition, 11 SSR markers were designed from the 184 kb contig containing the *Rps/k* region (GenBank accession no. EU450800), three of which (Satt1k2a, Satt1k3 and Satt1k4b) were polymorphic between ‘Yudou 29’ and ‘Jikedou 2’ (Supplementary Table 1).

Molecular analysis of the 214 $F_{2:3}$ families using these 11 polymorphic markers revealed that their segregation

Table 2 Genetic segregation of resistance to *P. sojae* isolate PsMC1 in the 214 $F_{2:3}$ families derived from the cross between Yudou 29 and Jikedou 2

| Cultivar and the cross | Generation | Amount | Observed number | | | Expected ratio and goodness of fit | | |
|------------------------|------------------|--------|-----------------|-----|----|------------------------------------|----------|----------|
| | | | R | Rs | S | (R:Rs:S) | χ^2 | <i>p</i> |
| Yudou 29 | P ₁ | 20 | 20 | – | – | | | |
| Jikedou 2 | P ₂ | 20 | – | – | 20 | | | |
| Jikedou 2 × Yudou 29 | F _{2:3} | 214 | 51 | 103 | 60 | 1:2:1 | 1.06 | 0.62 |

Fig. 2 Simple sequence repeat (SSR)-based genetic linkage map of the *RpsYD29* region. The map was deduced from segregation analysis of 214 $F_{2,3}$ families from a cross between *Glycine max* cultivars ‘Yudou 29’ and ‘Jikedou 2’. The map was generated in Joinmap using Kosambi’s mapping function. Marker names are on the right and distances (cM) on the left



pattern fits the 1:2:1 ratio (Supplementary Table 2). *RpsYD29* was closely linked to the SSR markers SattWM82-50 and Satt1k4b at a distance of 0.5 and 0.2 cM, respectively, and a genetic map consisting of the 11 SSR markers and *RpsYD29* was constructed (Fig. 2).

Candidate gene prediction

A BLAST search showed that the physical distance of the region between markers SattWM82-50 and Satt1k4b, which are at nucleotide positions 3,857,715 and 4,062,474, respectively, is approximately 204.8 kb. According to the Glyma1.0 annotations, a total of 11 genes have been identified in this region, of which 9 are supported by ESTs and two are putative (Supplementary Table 3).

Nucleotide binding site-leucine rich repeat (NBS-LRR) type resistance genes have been implicated in the resistance of *RpsIk* (Gao and Bhattacharyya 2008). Therefore, the identified 204.8 kb region containing *RpsYD29* was strictly evaluated for this motif, and two full-length NBS-LRR type genes, Glyma03g04030.1 and Glyma03g04080.1, were found to share amino acid sequence identities of 62–75 % with *Rps1k-1* and *Rps1k-2*.

Discussion

The soybean originated in China and large quantities of Chinese germplasms have been screened for PRR resistance (Tang et al. 2010; Xia et al. 2011; Zhu et al. 2006). All identified *Rps* genes (except *Rps8*) and *Rps* gene combinations are present in germplasms from China (Kyle et al. 1998; Lohnes

et al. 1994, 1996) and many germplasms are postulated to carry novel *Rps* genes (Xia et al. 2011; Zhu et al. 2006).

In this study, we investigated the resistance phenotype and genomic location of the *RpsYD29* gene in ‘Yudou 29’. The results confirm that *RpsYD29* is a strong, broad-spectrum resistance gene that is likely a novel allele at the *RpsI* locus, or a novel gene at another locus tightly linked to *RpsI* locus.

In addition to *RpsYD29*, eight known *Rps* genes were previously mapped to MLG N, including five alleles at the *RpsI* locus (Bernard et al. 1957; Buzzell and Anderson 1992; Mueller et al. 1978), *Rps7* (Weng et al. 2001), *RpsYD25* (*Rpsyu25*) (Fan et al. 2009; Sun et al. 2011) and an *Rps* gene in cv. Waseshiroge (Sugimoto et al. 2011). In our cluster analysis based on the Phytophthora resistance to 25 *P. sojae* isolates, ‘Yudou 29’ and the differentials carrying *RpsI* alleles (*Ia*, *Ib*, *Ic*, *Id*, *Ik*), *Rps7* and *RpsYD25* fell into different subgroups cutting at the coefficient 0.726, and ‘Yudou 29’ was clustered in a subgroup different from all these genes (except *RpsId*) on MLG N even cutting at 0.518. Sugimoto et al. (2008) stated that *RpsId* was located “above” Sat_186, and was flanked by Satt152 (11.5 cM) and Sat_186 (5.7 cM). In the present study, *RpsYD29* was located “below” Sat_186, and was flanked by Sat_186 (2.9 cM) and Satt530 (2.5 cM). These indicated that *RpsYD29* is a distinct gene from the *RpsI* alleles, *Rps7* and *RpsYD25*. In addition, the *Rps* gene in Waseshiroge was located between Satt009 and T003044871 and may reside in the nucleotide region between 3,919,203 and 4,486,048 of the chromosome 03 (MLG N) (Sugimoto et al. 2011). *RpsYD29* is not at the same location as the *Rps* gene in Waseshiroge and thus is distinct from this gene as well.

Carrying the novel resistance gene, *RpsYD29*, conveying broad-spectrum resistance to *P. sojae* isolates, ‘Yudou 29’ could be used as a new source of resistance for breeding Phytophthora-resistant cultivars. The markers identified in this study will facilitate the tracking of *RpsYD29* in progenies or cultivars for reliable MAS in breeding programs. The SSR markers SattWM82-50 and Satt1k4b positioned *RpsYD29* within very tight genetic distances of 0.5 and 0.2 cM, respectively. The accuracy of genotyping was theoretically estimated to be 99.60 and 99.80 % for MAS with SattWM82-50 and Satt1k4b, respectively. The selection accuracy would be as high as 99.99 % for MAS using both markers.

Acknowledgments The work was supported by the Special Fund for Agro-scientific Research in the Public Interest (3-20, 201303018) and the Program of Protection of Crop Germplasm Resources (NB2010-2130135-25-14) from the Ministry of Agriculture of the People’s Republic of China. We thank Dr. Schmitthener at the Ohio State University, USA, for providing the cultivars/lines carrying a single *Rps* gene at the *RpsI* to *Rps8* loci and the susceptible control cultivar Williams.

References

- Allen GC, Flores-Vergara MA, Krasynanski S, Kumar S, Thompson WF (2006) A modified protocol for rapid DNA isolation from plant tissues using cetyltrimethylammonium bromide. *Nat Protoc* 1:2320–2325
- Bernard RL, Cromeens CR (1981) An allele at the *Rps1* locus from the variety Kingwa. *Soybean Genet Newsl* 8:40–42
- Bernard RL, Smith PE, Kaufmann MJ, Schmitthenner AF (1957) Inheritance of resistance to *Phytophthora* root rot and stem rot in the soybean. *Agron J* 49:391
- Buzzell RI, Anderson TR (1992) Inheritance and race reaction of a new soybean *Rps1* allele. *Plant Dis* 76:600–601
- Chen QH, Wang QY, Wang YC, Zheng XB (2004) Identification and sequencing of ribosomal DNA-ITS of *Phytophthora sojae* in Fujian. *Acta Phytopathol Sinica* 34:112–116
- Chen XL, Zhu ZD, Wang XM, Xiao YN, Wu XF (2008) Postulation of *Phytophthora* resistance genes in soybean cultivars or lines. *Sci Agric Sinica* 41:1227–1234
- Cui KL, Yin WX, Tang QH, Dong SM, Zheng XB, Zhang ZG, Wang YC (2010) Distribution, pathotypes, and metalaxyl sensitivity of *Phytophthora sojae* from Heilongjiang and Fujian provinces in China. *Plant Dis* 94:881–884
- Dorrance AE, McClure SA, St Martin SK (2003) Effect of partial resistance on *Phytophthora* stem rot incidence and yield of soybean in Ohio. *Plant Dis* 87:308–312
- Fan AY, Wang XM, Fang XP, Wu XF, Zhu ZD (2009) Molecular identification of *Phytophthora* resistance gene in soybean cultivar Yudou 25. *Acta Agron Sinica* 35:1844–1850
- Gao H, Bhattacharyya MK (2008) The soybean-*Phytophthora* resistance locus *Rps1-k* encompasses coiled coil-nucleotide binding-leucine rich repeat-like genes and repetitive sequences. *BMC Plant Biol* 8:1–14
- Gao H, Narayanan NN, Ellison L, Bhattacharyya MK (2005) Two classes of highly similar coiled coil-nucleotide binding-leucine rich repeat genes isolated from the *Rps1-k* locus encode *Phytophthora* resistance in soybean. *Mol Plant Microbe Interact* 18:1035–1045
- Gordon SG, St Martin SK, Dorrance AE (2006) *Rps8* maps to a resistance gene rich region on soybean molecular linkage group F. *Crop Sci* 46:168–173
- Grau CR, Dorrance AE, Bond J, Russin J (2004) Fungal diseases. In: Boerma HR, Specht JE (eds) *Soybeans: Improvement, production and uses*. Agronomy Monograph, 3rd edn. American Society of Agronomy, Madison, pp 679–763
- Haas JH, Buzzell RI (1976) New races 5 and 6 of *Phytophthora megasperma* var. *sojae* and differential reactions of soybean cultivars for races 1 and 6. *Phytopathology* 66:1361–1362
- Kuan TL, Erwin DC (1980) *Formae speciales* differentiation of *Phytophthora megasperma* isolates from soybeans and alfalfa. *Phytopathology* 70:333–338
- Kyle DE, Nickell CD, Nelson RL, Pedersen WL (1998) Response of soybean accessions from provinces in southern China to *Phytophthora sojae*. *Plant Dis* 82:555–559
- Lohnes DG, Schmitthenner AF, Nickell CD (1994) Incidence and distribution of *Phytophthora* resistance in new soybean germplasm from China. *Agronomy Abstracts* 217
- Lohnes DG, Nickell CD, Schmitthenner AF (1996) Origin of soybean alleles for *Phytophthora* resistance in China. *Crop Sci* 36:1689–1692
- Michelmore RW, Paran I, Kesseli RV (1991) Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA* 88:9828–9832
- Mueller EH, Athow KL, Laviolette FA (1978) Inheritance of resistance to four physiologic races of *Phytophthora megasperma* var. *sojae*. *Phytopathology* 68:1318–1322
- Ploper LD, Athow KL, Laviolette FA (1985) A new allele at *Rps3* locus for resistance to *Phytophthora megasperma* f. sp. *glycinea* in soybean. *Phytopathology* 75:690–694
- Rohlf F (2000) NTSYS-PC numerical taxonomy and multivariate analysis system ver 2.11L. Appl Biostat, New York
- Sandhu D, Schallock KG, Rivera-Velez N, Lundeen P, Cianzio S, Bhattacharyya MK (2005) Soybean *Phytophthora* resistance gene *Rps8* maps closely to the *Rps3* region. *J Hered* 96:536–541
- Schmitthenner AF (1999) *Phytophthora* rot of soybean. In: Hartman GL, Sinclair JB, Rupe JC (eds) *Compendium of soybean diseases*, 4th edn. The American Phytopathological Society Press, St Paul, pp 39–42
- Schmitthenner AF, Hobe M, Bhat RG (1994) *Phytophthora sojae* races in Ohio over a 10-year interval. *Plant Dis* 78:269–276
- Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J, Xu D, Hellsten U, May GD, Yu Y, Sakurai T, Umezawa T, Bhattacharyya MK, Sandhu D, Valliyodan B, Lindquist E, Peto M, Grant D, Shu S, Goodstein D, Barry K, Futrell-Griggs M, Ab-ernathy B, Du J, Tian Z, Zhu L, Gill N, Joshi T, Libault M, Sethuraman A, Zhang XC, Shinozaki K, Nguyen HT, Wing RA, Cregan PB, Specht J, Grimwood J, Rokhsar D, Stacey G, Shoemaker RC, Jackson SA (2010) Genome sequence of the palaeopolyploid soybean. *Nature* 463:178–183
- Shen CY, Su YC (1991) Discovery and preliminary studies of *Phytophthora megasperma* on soybean in China. *Acta Phytopathol Sinica* 21:298
- Slaminko TL, Bowen CR, Hartman GL (2010) Multi-Year evolution of commercial soybean cultivars for resistance to *Phytophthora sojae*. *Plant Dis* 94:368–371
- Sugimoto T, Yoshida S, Watanabe K, Aino M, Kanto T, Maekawa K, Irie K (2008) Identification of SSR markers linked to the *Phytophthora* resistance gene *Rps1-d* in soybean. *Plant Breed* 127:154–159
- Sugimoto T, Yoshida S, Kaga A, Hajika M, Watanabe K, Aino M, Tatsuda K, Yamamoto R, Matoh T, Walker DR, Biggs AR, Ishimoto M (2011) Genetic analysis and Identification of DNA markers linked to a novel *Phytophthora sojae* resistance gene in the Japanese soybean cultivar Waseshiroge. *Euphytica* 182:133–145
- Sugimoto T, Kato M, Yoshida S, Matsumoto I, Kobayashi T, Kaga A, Hajika M, Yamamoto R, Watanabe K, Aino M, Matoh T, David R, Walker DR, Biggs AR, Ishimoto M (2012) Pathogenic diversity of *Phytophthora sojae* and breeding strategies to develop *Phytophthora*-resistant soybeans. *Breed Sci* 61:511–522
- Sun S, Wu XL, Zhao JM, Wang YC, Tang QH, Yu DY, Gai JY, Xing H (2011) Characterization and mapping of *RpsYu25*, a novel resistance gene to *Phytophthora sojae*. *Plant Breed* 130:139–143
- Tang QH, Cui LK, Li DL, Dai TT, Yin WX, Dong SM, Xing H, Zheng XB, Wang YC (2010) Resistance evaluation of soybean germplasm from Huanghuai Valley to *Phytophthora* Root Rot. *Sci Agric Sinica* 43:2246–2252
- Tyler BM (2007) *Phytophthora sojae*: root rot pathogen of soybean and model oomycete. *Mol Plant Pathol* 8:1–8
- Van Ooijen JW (2006) JoinMap 4 Software for the calculation of genetic linkage maps in experimental populations. *Kyazma B.V.*, Wageningen
- Weng C, Yu K, Anderson TR, Poysa V (2001) Mapping genes conferring resistance to *Phytophthora* root rot of soybean, *Rps1a* and *Rps7*. *J Hered* 92:442–446
- Wrather JA, Stienstra WC, Koenning SR (2001) Soybean disease loss estimates for the United States from 1996 to 1998. *Can J Plant Pathol* 23:122–131

- Xia CJ, Zhang JQ, Wang XM, Wu XF, Liu ZX, Zhu ZD (2011) Analyses of resistance genes to *Phytophthora* root rot in soybean germplasm. *Chin J Oil Crop Sci* 33:396–401
- Yao HY, Wang XM, Wu XF, Xiao YN, Zhu ZD (2010) Molecular mapping of *Phytophthora* resistance gene in soybean cultivar Zaoshu18. *J Plant Genet Resour* 11:213–217
- Yu AL, Xu PF, Wang JS, Zhang SZ, Wu JJ, Li WB, Chen WY, Li NH, Fan SJ, Wang X, Jiang LY (2010) Genetic analysis and SSR mapping of gene resistance to *Phytophthora sojae* race 1 in soybean cv Suinong 10. *Chin J Oil Crop Sci* 32:462–466
- Zhang SZ, Xu PF, Wu JJ, Xue AG, Zhang JX, Li WB, Chen C, Chen WY, Lv HY (2010) Races of *Phytophthora sojae* and their virulences on soybean cultivars in Heilongjiang, China. *Plant Dis* 94:87–91
- Zhu ZD, Wang HB, Wang XM, Chang RZ, Wu XF (2003) Distribution and virulence diversity of *Phytophthora sojae* in China. *Sci Agric Sinica* 36:793–799
- Zhu ZD, Huo YL, Wang XM, Huang JB, Wu XF (2006) Screening for resistance sources to *Phytophthora* root rot in soybean. *J Plant Genet Resour* 7:24–30
- Zhu ZD, Huo YL, Wang XM, Huang JB, Wu XF (2007) Molecular identification of a novel *Phytophthora* resistance gene in soybean. *Acta Agron Sinica* 33:154–157