

Research article

Open Access

Use of genome sequence data in the design and testing of SSR markers for *Phytophthora* species

Leonardo Schena*¹, Linda Cardle² and David EL Cooke²

Address: ¹Department of "Gestione dei Sistemi Agrari e Forestali", Mediterranean University of Reggio Calabria, Località Feo di Vito, 89124 Reggio Calabria, Italy and ²Scottish Crop Research Institute, Invergowrie, Dundee, DD2 5DA, Scotland, UK

Email: Leonardo Schena* - lschena@unirc.it; Linda Cardle - Linda.Milne@scri.ac.uk; David EL Cooke - David.Cooke@scri.ac.uk

* Corresponding author

Published: 19 December 2008

Received: 6 June 2008

BMC Genomics 2008, 9:620 doi:10.1186/1471-2164-9-620

Accepted: 19 December 2008

This article is available from: <http://www.biomedcentral.com/1471-2164/9/620>

© 2008 Schena et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Microsatellites or single sequence repeats (SSRs) are a powerful choice of marker in the study of *Phytophthora* population biology, epidemiology, ecology, genetics and evolution. A strategy was tested in which the publicly available unigene datasets extracted from genome sequences of *P. infestans*, *P. sojae* and *P. ramorum* were mined for candidate SSR markers that could be applied to a wide range of *Phytophthora* species.

Results: A first approach, aimed at the identification of polymorphic SSR loci common to many *Phytophthora* species, yielded 171 reliable sequences containing 211 SSRs. Microsatellites were identified from 16 target species representing the breadth of diversity across the genus. Repeat number ranged from 3 to 16 with most having seven repeats or less and four being the most commonly found. Trinucleotide repeats such as (AAG)_n, (AGG)_n and (AGC)_n were the most common followed by pentanucleotide, tetranucleotide and dinucleotide repeats. A second approach was aimed at the identification of useful loci common to a restricted number of species more closely related to *P. sojae* (*P. alni*, *P. cambivora*, *P. europaea* and *P. fragariae*). This analysis yielded 10 trinucleotide and 2 tetranucleotide SSRs which were repeated 4, 5 or 6 times.

Conclusion: Key studies on inter- and intra-specific variation of selected microsatellites remain. Despite the screening of conserved gene coding regions, the sequence diversity between species was high and the identification of useful SSR loci applicable to anything other than the most closely related pairs of *Phytophthora* species was challenging. That said, many novel SSR loci for species other than the three 'source species' (*P. infestans*, *P. sojae* and *P. ramorum*) are reported, offering great potential for the investigation of *Phytophthora* populations. In addition to the presence of microsatellites, many of the amplified regions may represent useful molecular marker regions for other studies as they are highly variable and easily amplifiable from different *Phytophthora* species.

Background

The genus *Phytophthora*, with other Oomycetes, fall within the kingdom Stramenopila, which also includes golden-brown algae, diatoms, and brown algae such as kelp [1-4]. This genus stands out among the plant pathogens since a

significant number of the 80 or so described species continue to prove a threat to ecosystem stability and plant productivity on a global scale [5-8]. Despite the importance of *Phytophthora* species, studies of their molecular diversity have been limited by the power of the genetic

markers and difficulties in comparing results among laboratories. Accurate studies based on the analysis of mitochondrial and nuclear DNA have resulted in a consensus of the phylogenetic relationships within the genus with a grouping into 10 genetically related clades now accepted [2,3,9]. However, these studies were based on genes commonly conserved within a species and therefore unsuitable to characterize intraspecific variability. Other approaches to study intraspecific variability among *Phytophthora* species including RAPD-PCR and AFLP have proved valuable within a particular study but comparing results from one laboratory to another has always proved challenging with such fingerprinting tools [10-13]. Although microsatellites or simple sequence repeats (SSRs) have been recognised as one of the most powerful choices of markers for molecular ecology they have only relatively recently been exploited in the study of *Phytophthora* populations. SSRs are tandemly repeated motifs of one to six bases which occur frequently and randomly in all eukaryotic genomes although their frequency varies significantly among different organisms [14]. They exhibit a high degree of length polymorphism among related organisms due to stepwise mutations affecting the number of repeat units and leading to polymorphism [14,15]. Dinucleotide repeats account for the majority of microsatellites for many species whereas trinucleotide and hexanucleotide repeats are the most likely repeat classes to appear in coding regions because they do not cause a frameshift [16,17]. Major advantages SSRs include: (i) multiple SSR alleles may be detected at a single locus using a simple PCR-based screen, (ii) SSRs are evenly distributed across the genome, (iii) they are co-dominant, (iv) very small quantities of DNA are required for screening, (v) analysis may be semi-automated, and (vi) results are objective compared to random amplification methods [18].

Microsatellites have been used to investigate genetic structure and reproductive biology of Oomycetes species including *Plasmopara viticola*, *P. cinnamomi*, *P. infestans*, and *P. ramorum* [19-21,23-25]. However, a major limitation to their wider exploitation is the need for prior species-specific marker isolation that requires knowledge of the DNA sequence of the SSR flanking regions to which specific primers have to be designed. Such regions are usually conserved within a species but the likelihood of primers successfully working between species decreases with increasing genetic distance and, in practice, primers are usually developed anew for each species [25,26]. Common methods for the discovery of SSR loci are based on constructing genomic DNA libraries enriched for SSR sequences. These methods were utilised for *P. cinnamomi* and *P. ramorum*, however they are time-consuming, and the specific sequencing of DNA libraries required is expensive [20,25]. Many commercial and academic laboratories

specialise in microsatellite isolation services and can provide a set of polymorphic microsatellite loci for a new species in 3–6 months for a cost of approximately USD 1,500 per locus, or USD 10,000 for 10–15 loci [14].

The availability of entire genome sequences for an increasing number of species including *P. infestans* <http://www.broad.mit.edu/>, *P. ramorum* and *P. sojae* <http://genome.jgi-psf.org/> have proved novel opportunities to identify and evaluate potential SSR markers identified by computational tools (Abajian, 1994, <http://espressosoft.ware.com/pages/sputnik.jsp>) [27,28]. This approach has been utilised to identify SSRs for the study of European and USA populations of *P. ramorum* and for monitoring the genetic variation in populations of *P. infestans* across Europe and worldwide [23,24,29,30].

Recently, Garnica et al. used an *in silico* approach to survey and compare simple sequence repeats (SSRs) in transcript sequences from the genomes of *P. sojae*, *P. ramorum* and *P. infestans* [27]. They also evaluated *in silico* transferability of SSRs among the *Phytophthora* species and found that a proportion (7.5%) of primers could, in theory, be transferred between at least two of the three species. In the present study SSRs from *P. infestans*, *P. sojae* and *P. ramorum* were analysed to identify useful loci common to many *Phytophthora* species (Approach 1) or to a restricted number of species closely related to *P. sojae* (Approach 2). Selected loci were amplified and sequenced from 16 (Approach 1) and 5 (Approach 2) different *Phytophthora* species and a comprehensive SSRs dataset was created.

Results

Approach 1 – SSRs for many *Phytophthora* species

The aim of this approach was to identify loci containing SSRs common to a large number of *Phytophthora* species (Fig. 1A). The method was validated using 16 different species (Table 1) representing the breadth of diversity across the genus [2,3].

Analysis of sequences from *P. infestans*, *P. sojae* and *P. ramorum* genome projects and scanning for homologous SSRs

Predicted gene datasets from *P. infestans* <http://www.broad.mit.edu/>, *P. sojae* and *P. ramorum* <http://genome.jgi-psf.org/> were scanned for the presence of microsatellites defined as short tandem repeat motifs (SSRs) of 2–6 bp. Both perfect and compound SSRs were selected with a minimal acceptable length of 10 bp (dinucleotide motifs) and 12 bp (tri- and tetranucleotide motifs). SSRs with a minimum of three repeats were included in the analyses of penta-nucleotide repeats. This search yielded 9333 sequences containing SSRs (1465 from *P. infestans*, 5348 from *P. sojae* and 2520 from *P. ramorum*). The relative abundance of SSRs was 103, 183 and 114 per Mb of

Table 1: Isolates of *Phytophthora* included in the study, their designations and origins.

<i>Phytophthora</i> species	Isolate numbers	Origin		
		Host	Country	Year
<i>P. alni</i> subsp. <i>alni</i>	SCR2P	<i>Alnus</i> sp.	UK	1995
	SCR2P4 ^(a)	<i>Alnus</i> sp.	Germany	1995
	SCR2P8 ^(a)	<i>Alnus</i> sp.	France	1996
<i>P. cambivora</i>	SCR2P67 (IMI 296831)	<i>Rubus idaeus</i>	Scotland	1985
	SCR2P75 ^(a)	<i>Fagus</i> sp.	UK	1995
	SCR2P80 ^(a)	<i>Castanea sativa</i>	Italy	1995
<i>P. cinnamomi</i>	SCR2P82 ^(a)	Eucalypt	Australia	
	SCR2P115 (CBS270.55)	<i>Chamaecyparis lawsoniana</i>	Netherlands	1993
	SCR2P118 (CBS342.72)	<i>Persea gratissima</i>	California	1972
<i>P. citricola</i>	SCR2P121 ^(a)		Australia	
	SCR2P130	<i>Rubus idaeus</i>	Scotland	1986
	SCR2P136 ^(a)	Soil	UK	1995
	SCR2P140 ^(a)	<i>Taxus</i> sp.	UK	1995
<i>P. europaea</i>	SCR2P143 ^(a)	<i>Quercus robur</i>	Germany	1994
	SCR2P622	<i>Quercus robur</i>	Switzerland	1995
<i>P. fragariae</i> var. <i>rubi</i>	SCR2P333 (IMI355974)	<i>Rubus idaeus</i>	Scotland	1985
<i>P. ilicis</i>	SCR2P377	<i>Ilex aquilifolium</i>	UK	1995
	SCR2P379 ^(a)	<i>Ilex aquilifolium</i>	UK	
<i>P. infestans</i>	SC03.26.3.3	<i>Solanum tuberosum</i>	Scotland	2003
<i>P. inundata</i>	SCR2P644 (IMI389751)	<i>Salix</i> sp.	UK	1972
<i>P. lateralis</i>	SCR2P390 (IMI 040503)	<i>Chamaecyparis lawsoniana</i>	U.S.A.	1942
<i>P. nemorosa</i>	SCR2P910			
<i>P. pseudosyringae</i>	SCR2P674 (IMI390500)	<i>Malus pumila</i>	Italy	2001
	SCR2P734 ^(a)	<i>Fagus sylvatica</i>	Italy	2003
<i>P. psychrophila</i>	SCR2P630	<i>Quercus ilex</i>	France	1996
<i>P. quercina</i>	SCR2P541	<i>Quercus robur</i>	Germany	1995
<i>P. ramorum</i>	SCR2P911	<i>Rhododendron</i> sp.	Scotland	2004
<i>P. sojae</i>	SCR2P555	<i>Glycine max</i>	USA	

^(a) = Additional isolates utilised to evaluate intraspecific variability

predicted gene sequence for *P. infestans*, *P. sojae* and *P. ramorum*, respectively.

Selected regions were compared by BLAST analysis to identify homologous regions flanking SSRs in at least two of the three species (*P. sojae*, *P. ramorum* and *P. infestans*). This analysis identified 4135 SSRs from *P. infestans* (688), *P. ramorum* (1470), and *P. sojae* (1977). A very limited number of loci containing SSRs were common to the three species; most loci were common to *P. ramorum* and *P. sojae* (81.6%), *P. infestans* and *P. ramorum* (7%) or *P. infestans* and *P. sojae* (11%). In most of the cases, homologous loci contained the same SSR motif in different *Phytophthora*s, however the number of repeats was consistently higher in the 'source' species than the other two.

Among the selected loci, the number of SSR repeats ranged from 3 to 13, from 3 to 12 and from 3 to 17 in *P. infestans*, *P. ramorum* and *P. sojae* respectively. Most SSRs showed seven repeats or less (98.2% *P. infestans*, 97.4% *P.*

ramorum, 94.4% *P. sojae*), with a repeat number of four being the most common in all species.

Selection and amplification of target regions containing SSRs

The 4135 homologous regions previously identified were manually analysed to select those with the highest number of repeats and flanked by the most conserved sequences on both sides. The latter condition was necessary to design primers suitable for as many species as possible. Based on this analysis 6, 7 and 12 target regions were identified across the genome of *P. infestans*, *P. ramorum* and *P. sojae* respectively. These regions, containing 8, 17 and 33 SSRs respectively, were selected (Table 2) for amplification from 16 different species of *Phytophthora* representing the breadth of diversity in the genus (Table 1). To this aim, a total number of 62 different degenerate primers (12 for *P. infestans*, 18 for *P. ramorum*, and 32 for *P. sojae*) were designed (Table 2). When target regions contained two or more SSRs and/or were too long to be amplified by a single amplification, a pool of different primers was designed (Table 2). Considerable effort was




Figure 1 Schematic representation of two different approaches utilised to identify SSRs in a broad range of *Phytophthora* species (A) or in *Phytophthora* spp. Clade 7 (B).

made to obtain successful amplification from as many species as possible. This involved screening of several primer pairs for each genomic region and for each *Phytophthora* species (Table 2) and adjustment of annealing

temperatures and MgCl₂ concentration for PCR reactions (Tables 3, 4, 5).

Table 2: Set of primers designed with the 1st approach (Fig. 1A) to amplify genomic regions with candidate SSRs in a broad range of *Phytophthora* species (Table 1).

	Forward primers ^(a)	Reverse primers ^(a)	SSRs ^(a)	Source ^(b)
<i>P. sojae</i>	S1F ACGACGTGTCCAAGAACCAC	S3R ATGTTGACCGTGTCTGCTG	(CCG) ₇ ; (AGC) ₄ ; (AGC) ₁₄	scaffold_3: 1346836–1347814
	S4F AARATGACGTGGACKGAGAG	S5R TGATSGTGGAGAARCTCATCT	(AAC) ₁₄	scaffold_14: 861958–862585
	S6F GGAGTTCGCCATCAACAAC	S7R TCAGCTTCTGTCGRTCGAC	(AAG) ₁₄	scaffold_24: 159790–160289
	S8F YGYGTCTCGCCCAYGAC	S9R GACGACACCGGSGAGAG	(ACC) ₄ ; (AGC) ₄ ; (AGC) ₂₈ ; (AAC) ₄	scaffold_76: 171071–172890
	S10F GCGSTACGAGACCTGGAC	S11R GACTCRCCCTTCGACTCSTC	(CAG) ₁₄	
	S12F GGAGGCCGAGTCGGARTA	S13R TAYTCCGACTCGGCCTCC	(AGC) ₁₄	scaffold_136: 56988–57479
	S14F GACGCMSSYYGAGTGAAAG	S15R ATTTKGSACAGATACCGACG	(AAG) ₁₅	scaffold_65: 332082–332852
	S16F TCTACGTGAATGCCATGAGG	S17R CGTTCAGCTTCTGTCGATCR	(AAG) ₁₅	scaffold_79: 50973–51428
	S18F YACCATCTCCAACCTGCTG	S19R CACCACCTCGAGTAGCTCCC	(AGC) ₇ ; (AGG) ₁₃	scaffold_2: 1159730–1160958
	S19F GGGAGCTACTCGAGGTGGTG	S20R TCGTCTCAATCTCKGACTGA	(AGC) ₆	

Table 2: Set of primers designed with the 1st approach (Fig. 1A) to amplify genomic regions with candidate SSRs in a broad range of *Phytophthora* species (Table 1). (Continued)

	S21F ATCTGGGCTCCASGAGGT	S22R CTGATCCTCCGCCACAY	(AAG) ₁₂ ; (ATC) ₆ ; (ATC) ₄ ; (AAG) ₅	scaffold_138: 18289–18854
	S23F GACTCGGACTCGGACGAC	S25R CTCCTGCTCKTCTTTCAGGC	(AGG) ₇ ; (AAG) ₁₀ ; (GAG) ₄ ; (AAG) ₁₂ ; (GAG) ₅ ; (AGA) ₆	scaffold_90: 249340–250137
		S37R CTTRCCBTCCTTGTCTTYT		
	S27F GAAGCGCGGGCGWGT	S31R TCCTCCTCTTCTTTCGTCW	(AAG) ₄ ; (AGG) ₄ ; (ACG) ₄ ; (AGG) ₁₁	scaffold_16: 680126–681136
	S31F WGACGAAGAAGAAGAGGAGGA	S28R TCATTCATCAGCGTGTCRAT	(GAG) ₄	
	S34F ABGAWGACGABGAGGAVGAV			
	S29F MGCAAGAAGGCGTCGTA	S30R CCTTCATCATGAGCTTCTGG	(AGG) ₄ (AAG) ₁₁	scaffold_40: 353572–354360
<i>P. ramorum</i>	R1F GYGGCGGTGGCTACAGYG	R3R CTGCTGYTGCTGGTTGAAAG	(ACC) ₄ ; (ACC) ₅ ; (ACC) ₄	scaffold_23: 349447–350425
	R2F CTACTCSAGCCGCTACGC			
	R3F CTTTCAACCAGCARGCAGCAG	R4R GTTTCATCATGCCWCCCATR	(AGC) ₈	scaffold_23: 350406–351828
	R4F YATGGGWGGCATGATGAAC	R5R AGGACCAGGAGATGGAGGAC	(AGC) ₄ ; (AGC) ₁₂ ; (AGC) ₄	
	R7F TGTTCCARACCCGCTTCC	R8R CACCAAGCAGCACKCGC	(ACG) ₉ ; (AAC) ₅ ; (AGC) ₁₀	scaffold_10: 436897–437690
		R9R GGAACGCACCAAAGACGC		

Table 2: Set of primers designed with the 1st approach (Fig. 1A) to amplify genomic regions with candidate SSRs in a broad range of *Phytophthora* species (Table 1). (Continued)

	R10F GGAGATGACGGAAGATGACG	R11R CCATCGAARTACATSACACGA	(AAGCC) ₄ ; (AGG) ₉ ; (AAG) ₇	scaffold_5: 750031-750612
	R13F AAGTCGAAGCTCGTGGTSAC	R14R GTATCCGCTGRAAGAGCGTC	(AAG) ₁₀	scaffold_78: 40203-40815
	R15F CCGGAGCGCGTGGGA	R16R GGTAGTTGAGCGGCTTCTTG	(CCG) ₆	scaffold_2778: 35-305
	R16F CAAGAAGCCGCTCAACTACC	R17R TAACGGATCAGCTCTTGCTG	(ATC) ₄ ; (AGG) ₈	scaffold_2778: 286-1134
<i>P. infestans</i>	I3F GCCTGTGGAYGAGAATGGYS	I4R CAGATCCACGACACCRGGY	(AAG) ₈	Pi_002_41652 _Feb05
	I5F CATCAACAAGTGCTCGTWCS	I6R TAGTCRAYGTTCTTGTTGTTCA	(AGC) ₅ ; (AGC) ₈	Supercont1.7 842678-843649
	I7F GHGTGGGCGAGTACTCCAAG	I8R AAGCTGGCTATRWACACTGCCG	(AG) ₉	Supercont1.4 1481811-1482015
	I9F GCATYGGGTCGTTCTCTGTA	I10R AGHGTGCAGTACAGACCCGC	(AAG) ₁₁	Supercont1.5 1235771-1236143
	I11F TCGTCBGTGTCCTCBACGTC	I12R ACCAGCATCTRTTCTGRGCAG	(ACC) ₈	Supercont1.45 522394-522633
	I13F GTCTGCGCTGTCGGAAC	I14R TRATGATGCGGTTTCATCTCG	(AAG) ₇ ; (AAG) ₄	Supercont1.220 167896-168460

(^a) = Primers are listed according to the localization in their respective genome projects (*P. sojae*, *P. ramorum* or *P. infestans*) and according to the flanked SSRs. In some circumstances, a pool of different primers was designed to amplify selected genomic regions from as many as possible *Phytophthora* species. Similarly, when target regions contained two or more SSRs and were too long to be amplified by a single amplification a pool of different primers was designed.

(^b) = Gene sequences available at <http://genome.jgi-psf.org/sojae1/sojae1.home.html> (*P. sojae*), <http://genome.jgi-psf.org/ramorum1/ramorum1.home.html> (*P. ramorum*) and <http://www.pfgd.org/> or http://www.broad.mit.edu/annotation/genome/phytophthora_infestans/Home.html (*P. infestans*).

Table 3: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. sojae* with the first approach (Fig. 1A).

Phytoph. species	SELECTED PRIMERS ^(a)											
	S1F-S3R	S4F-S5R	S6F-S7R	S10F-11R	S16F-17R	S18F-19R	S19F-20R	S21F-22R	S23F-S25R ⁽¹⁾	⁽²⁾ S31F-28R	S27F-S31R	S29F-S30R
<i>P. alni</i> subsp. <i>alni</i> SCR2P2	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216617</u> No SSR 58°C/1.0 mM*	<u>EF216607</u> (aac) ₄ 58°C/1.7 mM*	<u>EF216602</u> (agc) ₆ 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	<u>EF216590</u> No SSR 58°C/1.0 mM*	<u>EF216580</u> (aag) ₅ (agg) ₅ (aag) ₄ (aag) ₆ (agg) ₅ 58°C/1.0 mM*	<u>EF216565</u> (agg) ₇ (aag) ₁₀ (agg) ₁₆ (aag) ₁₂ (aag) ₄ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216555</u> (aag) ₈ (agg) ₄ (agg) ₄ (aag) ₅ (agg) ₄ (aag) ₄ (aag) ₅ 58°C/1.0 mM*	<u>EF216552</u> (aag) ₈ 58°C/1.0 mM*
<i>P. cambiv.</i> SCR67	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216618</u> No SSR 58°C/1.0 mM*	<u>EF216606</u> (aac) ₄ 58°C/1.7 mM*	<u>EF216601</u> (agc) ₄ (cg) ₅ 58°C/1.0 mM*	<u>EF216593</u> No SSR 55°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216581</u> (agg) ₄ (aag) ₅ (aag) ₄ 58°C/1.0 mM*	<u>EF216569</u> (agg) ₁₀ (aag) ₁₀ (agg) ₉ (aag) ₁₂ (agg) ₄ (agg) ₇ 58°C/1.0 mM*	NA ⁺	NA ⁺	<u>EF216551</u> (aag) ₈ 58°C/1.0 mM*
<i>P. cinnam.</i> SCR115	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	<u>EF2165</u> (aag) ₈ (agg) ₅ (aag) ₉ 58°C/1.0 mM*
<i>P. citricola</i> SCR130	NA ⁺	<u>EF216619</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216553</u> (aag) ₇ (aag) ₇ 58°C/1.0 mM*

Table 3: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. sojae* with the first approach (Fig. 1A). (Continued)

<i>P. europaea</i> SCR622	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216616</u> (acg) ₄ 58°C/1.0 mM*	<u>EF216605</u> No SSR 58°C/1.7 mM*	<u>EF216600</u> No SSR 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	NS ⁺⁺ 55°C/1.7 mM*	<u>EF216589</u> No SSR 58°C/1.0 mM*	<u>EF216576</u> (aag) ₄ 58°C/1.0 mM*	<u>EF216568</u> (agg) ₉ (aag) ₁₀ (aag) ₇ (agg) ₄ (aag) ₆ (agg) ₄ (agg) ₄ 58°C/1.0 mM*	NA ⁺	NA ⁺	<u>EF21655054</u> (aag) ₉ 58°C/1.0 mM*
<i>P. fragariae</i> <i>var. rubi</i> SCR333	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216594</u> (aac) ₄ 55°C/1.7 mM*	NA ⁺	<u>EF216584</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216542</u> (aag) ₇ (aag) ₈ 58°C/1.0 mM*
<i>P. ilicis</i> SCR377	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216608</u> (ccg) ₄ 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	NA ⁺	NA ⁺	<u>EF216575</u> No SSR 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216543</u> (aag) ₅ 58°C/1.0 mM*
<i>P. infestans</i> sc 03.26.3.3	NA ⁺	<u>EF216615</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	<u>EF216560</u> No SSR 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*
<i>P. inundata</i> SCR644	<u>EF216624</u> No SSR 58°C/1.0 mM*	<u>EF216614</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216599</u> No SSR 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	<u>EF216588</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/0.7 mM*	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216549</u> (aag) ₈ 58°C/1.0 mM*
<i>P. lateralis</i> SCR390	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216611</u> No SSR 58°C/1.0 mM*	<u>EF216604</u> (ac) ₅ 58°C/1.7 mM*	<u>EF216598</u> (agc) ₄ 58°C/1.0 mM*	NS ⁺⁺ 55°C/1.7 mM*	<u>EF216592</u> (acg) ₄ (agc) ₄ 58°C/1.0 mM*	<u>EF216587</u> No SSR 58°C/1.0 mM*	<u>EF216579</u> (aag) ₅ 58°C/1.0 mM*	<u>EF216564</u> (agg) ₅ (aag) ₄ (aag) ₆ (agg) ₄ 58°C/1.0 mM*	<u>EF216556</u> (agg) ₄ 58°C/0.7 mM*	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216548</u> (aag) ₅ (aag) ₉ 58°C/1.0 mM*
<i>P. nemor.</i> SCR910	NA ⁺	<u>EF216613</u> (ccg) ₄ 58°C/1.0 mM*	NA ⁺	<u>EF216597</u> No SSR 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 55°C/1.7 mM*	NA ⁺	NA ⁺	<u>EF216572</u> No SSR 58°C/1.0 mM*	<u>EF216559</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216547</u> (aag) ₅ 58°C/1.0 mM*

Table 3: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. sojae* with the first approach (Fig. 1A). (Continued)

<i>P. pseudos.</i> SCR674	<u>EF216622</u> No SSR 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216596</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	<u>EF216573</u> No SSR 58°C/1.0 mM*	<u>EF216557</u> (agg) ₁₄ 58°C/1.0 mM*	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216546</u> (aag) ₄ 58°C/1.0 mM*
<i>P. psychro.</i> SCR630	<u>EF216623</u> (agc) ₄ 58°C/1.0 mM*	<u>EF216612</u> (ccg) ₄ 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216574</u> No SSR 58°C/1.0 mM*	<u>EF216558</u> (agg) ₆ (agg) ₄ (aag) ₄ (agg) ₄ 58°C/0.7 mM*	NS ⁺⁺ 60°C/0.7 mM*	<u>EF216545</u> (aag) ₄ 58°C/1.0 mM*
<i>P. quercina</i> SCR6541	<u>EF216621</u> (agc) ₅ 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216563</u> (aag) ₁₀ (aag) ₅ 58°C/1.0 mM*	<u>EF216561</u> (agg) ₄ 58°C/1.0 mM*	NA ⁺	<u>EF216544</u> (aag) ₇ 58°C/1.0 mM*
<i>P. ramorum</i> SCR6911	<u>EF216620</u> No SSR 58°C/1.0 mM*	<u>EF216610</u> (acc) ₄ 58°C/1.0 mM*	<u>EF216603</u> No SSR 55°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216595</u> No SSR 55°C/1.7 mM*	<u>EF216591</u> No SSR 58°C/1.0 mM*	<u>EF216586</u> No SSR 58°C/1.0 mM*	<u>EF216578</u> (aag) ₄ (atc) ₄ (aag) ₅ (aag) ₅ 58°C/1.0 mM*	<u>EF216562</u> (aag) ₄ (agg) ₄ (aag) ₇ 58°C/1.0 mM*	NS ⁺⁺ 58°C/0.7 mM*	NA ⁺	<u>EF216540</u> (aag) ₄ 58°C/1.0 mM*
<i>P. sojae</i> SCR6555	NA ⁺	<u>EF216609</u> (aac) ₁₄ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	<u>EF382779</u> No SSR 55°C/1.7 mM*	NA ⁺	<u>EF216585</u> (agc) ₆ 58°C/1.0 mM*	<u>EF216577</u> (aag) ₁₂ (atc) ₆ (atc) ₄ (aag) ₅ 58°C/1.0 mM*	NS ⁺⁺ 58°C/0.7 mM*	NS ⁺⁺ 58°C/0.7 mM*	NA ⁺	<u>EF216541</u> (agg) ₄ (aag) ₁₁ 58°C/1.0 mM*

(a) = Primers listed in Table 2 and not reported in this table are those that did not produce any reliable sequence (see text).

(1) = Primer S25R was replaced by primer S37R for *P. nemorosa*, *P. pseudosyringae*, *P. psychrophila*, and *P. ilicis*.

(2) = Primer S31F was replaced by primer S34F for *P. alni*, *P. citricola*, *P. infestans*, *P. nemorosa*, and *P. quercina*.

NA⁺ = Isolate-primer combinations that did not produce any amplification or produced complex profiles (two or more fragments).

NS⁺⁺ = Isolate-primer combinations for which single PCR bands were obtained but direct sequencing did not produce reliable results.

°C/mM* = Selected annealing temperature (°C) and MgCl₂ concentration (mM) in PCR reactions.

Table 4: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. ramorum* with the first approach (Fig. 1A).

Phytophthora species	SELECTED PRIMERS ^(a)						
	⁽¹⁾ R1F-R3R	R3F-R4R	R4F-R5R	R7F-9R ⁽²⁾	R10F-R11R	R13F-R14R	R16F-R17R
<i>P. alni</i> subsp. <i>alni</i> SCR2P2	NA ⁺	NA ⁺	<u>EF216645</u> No SSR 58°C/1.0 mM*	<u>EF216671</u> No SSR 58°C/1.0 mM*	<u>EF216662</u> No SSR 58°C/1.7 mM*	<u>EF216651</u> (aag) ₄ 58°C/1.7 mM*	<u>EF216625</u> (acg) ₄ 58°C/1.0 mM*
<i>P. cambivora</i> SCR67	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216673</u> (agc) ₈ 58°C/1.0 mM*	<u>EF216663</u> (agg) ₄ (aag) ₄ (agg) ₄ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216626</u> (acg) ₄ 58°C/1.0 mM*
<i>P. cinnamomi</i> SCR115	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/ 1.7 mM*	NA ⁺
<i>P. citricola</i> SCR130	NA ⁺	NA ⁺	<u>EF216644</u> No SSR 58°C/1.0 mM*	NA ⁺	NA ⁺	NA ⁺	<u>EF216630</u> No SSR 58°C/1.0 mM*
<i>P. europaea</i> SCR622	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216643</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216661</u> (agg) ₄ (aag) ₇ (agg) ₅ 58°C/1.7 mM*	<u>EF216655</u> (aag) ₄ 58°C/1.7 mM*	NA ⁺
<i>P. fragariae</i> <i>var. rubi</i> SCR333	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.0 mM**	<u>EF216634</u> No SSR 58°C/1.0 mM*	<u>EF216672</u> (agc) ₇ 58°C/1.0 mM*	<u>EF216657</u> (aag) ₄ (agg) ₄ 58°C/1.7 mM*	<u>EF216647</u> (aag) ₄ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*
<i>P. ilicis</i> SCR377	NS ⁺⁺ 58°C/1.0 mM*	<u>EF216646</u> (agc) ₄ (agc) ₄ 58°C/1.0 mM*	<u>EF216635</u> (agc) ₄ (accat) ₅ 58°C/1.0 mM*	<u>EF216664</u> (actg) ₃ (agc) ₆ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216648</u> No SSR 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*
<i>P. infestans</i> sc 03.26.3.3	NS ⁺⁺ 58°C/ 1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NS ⁺⁺ 58°C/1.7 mM*	NA ⁺	NA ⁺
<i>P. inundata</i> SCR644	<u>EF216633</u> No SSR 58°C/ 1.0 mM*	NA ⁺	NA ⁺	NA ⁺	NA ⁺	<u>EF216654</u> No SSR 58°C/1.7 mM*	<u>EF216629</u> No SSR 58°C/1.0 mM*

Table 4: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. ramorum* with the first approach (Fig. 1A). (Continued)

<i>P. lateralis</i> SCR390	<u>EF216631</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216642</u> (aac) ₅ (agc) ₅ 58°C/1.7 mM*	<u>EF216669</u> (agc) ₅ (aac) ₆ 58°C/1.0 mM*	<u>EF216660</u> (agg) ₄ 58°C/1.7 mM*	NA ⁺	<u>EF216628</u> (agg) ₈ (agc) ₄ 58°C/1.0 mM*
<i>P. nemorosa</i> SCR910	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216639</u> (agc) ₄ (accat) ₄ 58°C/1.0 mM*	<u>EF216668</u> (actg) ₃ (agc) ₆ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216653</u> (agg) ₉ 58°C/1.7 mM*	NA ⁺
<i>P. pseudosyringae</i> SCR674	<u>EF216632</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216641</u> (accat) ₄ 58°C/1.0 mM*	<u>EF216667</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216652</u> No SSR 58°C/1.7 mM*	NA ⁺
<i>P. psychrophila</i> SCR630	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216640</u> (accat) ₄ 58°C/1.0 mM*	<u>EF216666</u> (actg) ₃ (agc) ₆ 58°C/1.0 mM*	NA ⁺	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*
<i>P. quercina</i> SCR541	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216638</u> No SSR 58°C/1.0 mM*	<u>EF216674</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216651</u> (aag) ₈ (aag) ₄ 58°C/1.7 mM*	NA ⁺
<i>P. ramorum</i> SCR911	NS ⁺⁺ 58°C/1.0 mM*	NA ⁺	<u>EF216637</u> (agc) ₄ (agc) ₁₀ (agc) ₄ 58°C/1.0 mM*	<u>EF216665</u> (agc) ₂₄ 58°C/1.0 mM*	<u>EF216659</u> (aagcc) ₄ (agg) ₉ (aag) ₇ 58°C/1.7 mM*	<u>EF216650</u> (aag) ₁₀ 58°C/1.7 mM*	<u>EF216627</u> (atc) ₄ (agg) ₈ 58°C/1.0 mM*
<i>P. sojae</i> SCR555	NA ⁺	NA ⁺	<u>EF216636</u> (agc) ₅ 58°C/1.0 mM*	<u>EF216670</u> (agc) ₁₀ (agcg) ₅ 58°C/1.0 mM*	<u>EF216658</u> (agg) ₅ (aag) ₆ (agg) ₄ 58°C/1.7 mM*	<u>EF216649</u> (agc) ₄ (aag) ₄ 58°C/1.7 mM*	NA ⁺

(a) = Primers listed in Table 2 and not reported in this table are those that did not produce any reliable sequence (see text).

(1) = Primer R1F was replaced by primer R2F for *P. cambivora*, *P. inundata*, *P. nemorosa*, *P. ilicis* and *P. fragaria*.

(2) = Primer R9R was replaced by primer R8R for *P. quercina*.

NA⁺ = Isolate-primer combinations that did not produce any amplification or produced complex profiles (two or more fragments).

NS⁺⁺ = Isolate-primer combinations for which single PCR bands were obtained but direct sequencing did not produce reliable results.

°C/mM* = Selected annealing temperature (°C) and MgCl₂ concentration (mM) in PCR reactions.

Table 5: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. infestans* with the first approach (Fig. 1A).

<i>Phytophthora</i> species	SELECTED PRIMERS ^(a)					
	I3F-4R	I5F-I6R	I7F-I8R	I9F-I10R	I11F-I12R	I13F-I14R
<i>P. alni</i> subsp. <i>alni</i> SCR2P	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216535</u> No SSR; 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216513</u> (aag) ₄ (agg) ₆ 58°C/1.0 mM*	NA ⁺	<u>EF216477</u> (agg) ₄ (aag) ₅ 58°C/1.7 mM*
<i>P. cambivora</i> SCR67	NS ⁺⁺ 58°C/1.7 mM*	NA ⁺	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216516</u> (acg) ₄ (aag) ₅ (agg) ₆ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216478</u> (agg) ₅ (aag) ₅ 58°C 1.7 mM*
<i>P. cinnamomi</i> SCRPI15	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216509</u> (aag) ₁₄ 58°C/1.0 mM*	<u>EF216494</u> (aag) ₄ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.7 mM*
<i>P. citricola</i> SCRPI30	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216534</u> (agc) ₄ ; (agc) ₅ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216520</u> (aag) ₄ 58°C/1.0 mM*	<u>EF216498</u> (aag) ₄ 58°C/1.7 mM*	<u>EF216482</u> (agg) ₉ (aag) ₅ 58°C/1.7 mM
<i>P. europaea</i> SCR622	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216512</u> No SSR 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216476</u> (agg) ₅ (aag) ₅ 58°C/1.7 mM*
<i>P. fragariae</i> var. <i>rubi</i> SCR333	NS ⁺⁺ 58°C/1.7 mM*	NA ⁺	NA ⁺	<u>EF216500</u> (acg) ₄ 58°C/1.0 mM*	NA ⁺	NA ⁺
<i>P. ilicis</i> SCR377	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216532</u> (agc) ₉ 58°C/1.0 mM*	NA ⁺	<u>EF216501</u> No SSR 58°C/1.0 mM*	<u>EF216495</u> No SSR 58°C/1.7 mM*	<u>EF216483</u> (aag) ₄ (agc) ₄ 58°C/1.7 mM
<i>P. infestans</i> sc 03.26.3.3	NS ⁺⁺ 58°C 1.7 mM*	<u>EF216524</u> (agc) ₆ (agc) ₅ 58°C/1.0 mM	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216499</u> (aag) ₁₁ 58°C/1.0 mM*	<u>EF216487</u> (acc) ₈ 58°C/1.7 mM*	<u>EF216474</u> (aag) ₇ (aag) ₄ 58°C/1.7 mM

Table 5: Accession numbers and SSRs for GenBank deposited sequences <http://www.ncbi.nlm.nih.gov/> amplified from 16 *Phytophthora* species (Table 1) using primers designed on *P. infestans* with the first approach (Fig. 1A). (Continued)

<i>P. inundata</i> SCR644	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216508</u> No SSR 58°C/1.0 mM*	<u>EF216497</u> No SSR 58°C/1.7 mM*	NA ⁺
<i>P. lateralis</i> SCR6390	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216527</u> (agc) ₄ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216507</u> No SSR 58°C/1.0 mM*	<u>EF216493</u> No SSR 58°C/1.7 mM*	<u>EF216481</u> (agg) ₅ (aag) ₄ 58°C/1.7 mM*
<i>P. nemorosa</i> SCR6910	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216531</u> (agc) ₇ 58°C/1.0 mM*	NA ⁺	<u>EF216503</u> No SSR 58°C/1.0 mM*	<u>EF216492</u> (aag) ₄ 58°C/1.7 mM*	<u>EF216486</u> (agc) ₄ (aag) ₄ (agc) ₄ 58°C/1.7 mM*
<i>P. pseudosyringae</i> SCR674	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216529</u> (agc) ₇ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216502</u> No SSR 58°C/1.0 mM*	<u>EF216496</u> (aag) ₅ 58°C/1.7 mM*	<u>EF216485</u> (agc) ₄ (aag) ₄ 58°C/1.7 mM*
<i>P. psychrophila</i> SCR630	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216528</u> (agc) ₄ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	NA ⁺	<u>EF216491</u> (aag) ₄ 58°C/1.7 mM*	<u>EF216484</u> (aag) ₄ (agc) ₄ 58°C/1.7 mM*
<i>P. quercina</i> SCR6541	NS ⁺⁺ 58°C/1.7 mM*	NS ⁺⁺ 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216506</u> No SSR 58°C/1.0 mM*	<u>EF216490</u> (aag) ₅ 58°C/1.7 mM*	<u>EF216480</u> (agg) ₆ (aag) ₄ 58°C/1.7 mM*
<i>P. ramorum</i> SCR6911	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216525</u> No SSR 58°C/1.0 mM*	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216505</u> No SSR 58°C/1.0 mM*	<u>EF216489</u> (acc) ₄ 58°C/1.7 mM*	<u>EF216479</u> (aac) ₇ (agg) ₉ 58°C/1.7 mM
<i>P. sojae</i> SCR6555	NS ⁺⁺ 58°C/1.7 mM*	<u>EF216526</u> No SSR 58°C/1.0 mM*	NA ⁺	<u>EF216504</u> (agc) ₄ 58°C/1.0 mM*	<u>EF216488</u> No SSR 58°C/1.7 mM*	<u>EF216475</u> (agg) ₇ (aag) ₅ 58°C/1.7 mM*

(a) = Primers listed in Table 2 and not reported in this table are those that did not produce any reliable sequence (see text).

NA⁺ = Isolate-primer combinations that did not produce any amplification or produced complex profiles (two or more fragments).

NS⁺⁺ = Isolate-primer combinations for which single PCR bands were obtained but direct sequencing did not produce reliable results.

°C/mM* = Selected annealing temperature (°C) and MgCl₂ concentration (mM) in PCR reactions.

The resultant primers enabled the amplification of 271 single PCR bands of the expected size (Fig. 2). In the remaining primer-species combinations, 193 amplifications did not produce any product or produced complex profiles (two or more PCR fragments) impeding direct sequencing (Fig. 2). Some primer combinations failed to amplify a product from any of the *Phytophthora* species whereas other combinations amplified single bands from all or most *Phytophthora* species (Tables 3, 4, 5).

Sequencing of single PCR bands and scanning for SSRs

All single PCR bands (271) were purified to remove excess primers and nucleotides and sequenced in both directions using the same primers used for the amplification. When forward and/or reverse sequences were not identical, amplification, purification and sequencing were repeated twice and all unreliable sequences were discarded. Finally, 171 sequences were obtained with primers designed




Figure 2
Amplification results obtained with 16 *Phytophthora* species (Table 1) using primers designed against *P. sojae*, *P. ramorum* and *P. infestans* genomes using Approach I (Fig. 1A). NA represents primer-species combinations in which amplification reactions did not produce any product or produced complex profiles (two or more PCR fragments) preventing direct sequencing. NS represents primer-species combinations in which amplification reactions produced single PCR bands, however direct sequencing did not yield reliable sequences. SQ represents primer-species combinations in which reliable sequences were obtained.

against *P. sojae* (70), *P. ramorum* (50) and *P. infestans* (51) genomes (Fig. 2) and scanned to identify SSRs by means of sputnik. Sequenced regions contained a total number of 211 SSRs distributed across the genome of the 16 target species with those of clade 7 (*P. alni*, *P. cambivora*, *P. europaea*, *P. fragariae* and *P. sojae*) and clade 8 (*P. lateralis* and *P. ramorum*) more highly represented (Fig. 3; Tables 3, 4, 5). A single microsatellite was identified in *P. inundata*. All SSRs identified in *P. infestans* were amplified with primers designed against its own genome (Fig. 3). Identified SSRs ranged in the number of repeats from 4 to 16, from 3 to 16 and from 4 to 14 in *P. sojae*, *P. ramorum* and *P. infestans* respectively (Tables 3, 4, 5). A single repeat of 24 was found in an SCRI isolate of *P. ramorum* (Table 4). Most SSRs were of seven repeats or less (88.9% *P. infestans*, 82.8% *P. ramorum*, 76.8% *P. sojae*), with a repeat number of four being the most common in all species (Fig. 4). Overall, the most common motifs were (AAG)_n, (AGG)_n and (AGC)_n representing 40.9%, 23.3% and 17.6% respectively of the total number of identified SSRs (Fig. 5). Trinucleotide repeats were the most common (94.7%) followed by pentanucleotide (2.4%), tetranucleotide (1.9%) and dinucleotide (1.0%) repeats.

To evaluate intraspecific variability a few selected target regions amplified by primers S23F-S25R, S21F-S22R, I9-I10 and I5-6 (Table 2) were examined and sequenced from additional isolates of *P. alni*, *P. cambivora*, *P. cinnamomi*, *P. pseudosyringae* and *P. ilicis* (Table 1). The analysed target regions did not show intraspecific variability among analysed isolates of *P. alni* subsp. *alni*, *P. pseudosyringae* or *P. ilicis*, whereas *P. cambivora* and *P. cinnamomi* isolates were polymorphic in all the tested primer combinations. As an example, the target region amplified with primers I9-I10 from *P. cinnamomi* was characterised by 12, 14 and 18 repeated motifs (AGG) in three tested isolates (Table 6).

Approach 2 – Identification of SSRs in *Phytophthora* spp. clade 7

The aim of this approach was to focus the search for SSR loci to a restricted range of four clade 7 *Phytophthora* species (*P. alni*, *P. cambivora*, *P. europaea* and *P. fragariae*) phylogenetically related to *P. sojae* (Fig 1B) [9].

Identification of target regions

This approach was based on a detailed list of SSRs identified in the genome of *P. sojae* and provided by Dr. Niklaus Grunwald at the Agricultural Research Service, U.S. Department of Agriculture, Corvallis, Oregon. Among the list, sixty genomic regions (500–1000 bp) were manually selected on the basis of having the longest SSRs in exons (20), introns (20) and non coding regions (20). The selected regions (Table 7) were screened using BLAST against the entire genomes of *P. ramorum* and *P. infestans*




Figure 3
 Number of SSRs identified for each of the 16 *Phytophthora* species using primers designed against *P. sojae*, *P. ramorum* and *P. infestans* genomes (Approach I, Fig. 1A).




Figure 4
Number of repeated motifs identified in 16 target *Phytophthora* species (Table 1) using primers designed against *P. sojae*, *P. ramorum* and *P. infestans* genomes according to Approach 1 (Fig. 1A).

to search for homology irrespective of the SSR regions. None of these regions aligned with sequences from the *P. infestans* genome whereas 18 of the 60 regions were sufficiently conserved to match homologous genes in *P. ramorum* (6 were localised in exons and 12 in introns). Surprisingly none of these 18 regions contained SSRs in *P. ramorum*, however it was hypothesised that microsatellites could be present in homologous regions of other *Phytophthora* species more closely related to *P. sojae*. To verify this hypothesis, thirty-six primers (18 pairs) were designed in the conserved flanking regions and used to amplify the target regions from *P. alni*, *P. cambivora*, *P. europaea*, *P. fragariae* and an SCRI isolate of *P. sojae* (Table 7). Degenerate primers were designed when necessary.

Amplification, sequencing and SSR scoring

Most primer-species combinations produced single PCR bands of the expected size (Table 8). Purification and direct sequencing of these PCR fragments produced 54 reliable sequences which were analysed as previously




Figure 5
List and frequency of the different SSR motifs identified in 16 *Phytophthora* species (Table 1) using primers designed on *P. sojae*, *P. ramorum* and *P. infestans* genomes according to Approach 1 (Fig. 1A).

described for Approach 1. Twelve different microsatellites were identified: 2 in *P. europaea*, 3 in *P. fragariae* and *P. alni* and 4 in *P. cambivora* (Table 8). Among these, 10 were trinucleotides and 2 were tetranucleotides repeated 4, 5 or 6 times. All regions sequenced from the SCRI isolate of *P. sojae* contained the predicted/expected SSR (Table 8).

Discussion

The present study was undertaken to develop a method to rapidly identify loci containing SSRs and to create a pool of microsatellite markers for species of the genus *Phytophthora* taking advantage of publicly available sequences for *P. sojae*, *P. ramorum* and *P. infestans*. Recently, Garnica et al. explored the transferability of microsatellites across *P. sojae*, *P. ramorum* and *P. infestans* via an *in silico* virtual PCR approach [27]. In the present study, such an analysis on the same three species was conducted but also followed up with a comprehensive screening and validation process on multiple species to provide a practical evaluation of

Table 6: Accession numbers and SSRs for selected microsatellites amplified and sequenced from two or more isolates of the same species to evaluate intraspecific variability.

<i>Phytophthora</i> species	<i>Phytophthora</i> isolates	Primers	SSRs	Accession number
<i>P. alni</i> subsp. <i>alni</i>	SCR2P	S23F-S25R	(agg) ₇ ; (aag) ₁₀ ; (agg) ₁₆ ; (aag) ₁₂ ; (aag) ₄	EF216565
	SCR4P		(agg) ₇ ; (aag) ₁₀ ; (agg) ₁₆ ; (aag) ₁₂ ; (aag) ₄	EF216567
	SCR8P		(agg) ₇ ; (aag) ₁₀ ; (agg) ₁₆ ; (aag) ₁₂ ; (aag) ₄	EF216566
<i>P. cambivora</i>	SCR67P	S23F-S25R	(agg) ₁₀ ; (aag) ₁₀ ; (agg) ₉ ; (aag) ₁₂ ; (agg) ₄ ; (agg) ₇	EF216569
	SCR75P		(agg) ₉ ; (aag) ₁₀ ; (agg) ₁₁ ; (aag) ₁₂ ; (agg) ₆	EF216571
	SCR82P		(agg) ₁₀ ; (aag) ₁₀ ; (agg) ₉ ; (aag) ₁₂ ; (agg) ₅	EF216570
<i>P. cambivora</i>	SCR67P	S21F-S22R	(agg) ₄ ; (aag) ₅ ; (aag) ₄	EF216581
	SCR80P		(agg) ₄ ; (aag) ₅ ; (aag) ₄	EF216582
	SCR82P		(aag) ₄ ; (agg) ₆ ; (agg) ₄	EF216583
<i>P. cambivora</i>	SCR67P	I9F-I10R	(acg) ₄ ; (aag) ₄ ; (agg) ₆	EF216516
	SCR75P		(acg) ₄ ; (aag) ₄ ; (agg) ₆	EF216519
	SCR80P		(acg) ₄ ; (aag) ₄ ; (agg) ₆	EF216517
	SCR82P		(aag) ₅ ; (agg) ₅	EF216518
<i>P. cinnamomi</i>	SCR115P	I9F-I10R	(aag) ₁₄	EF216509
	SCR118P		(aag) ₁₈	EF216511
	SCR121P		(aag) ₁₂	EF216510
<i>P. pseudosyringae</i>	SCR674P	I5F-I6R	(agc) ₇	EF216529
	SCR734P		(agc) ₇	EF216530
<i>P. ilicis</i>	SCR377P	I5F-I6R	(agc) ₉	EF216532
	SCR379P		(agc) ₉	EF216533

the procedure as a means of accelerating the search for new SSR markers in the genus *Phytophthora*.

The first approach was aimed at the identification of informative SSR loci common to many *Phytophthora* species. This approach was based on the hypothesis that among the large number of microsatellites distributed across the genome of species of the genus there may be a proportion in genes common to many species with sufficient sequence conservation in flanking regions to allow the design and use of universal SSR primers. Our search of the predicted gene sets yielded approximately 10% fewer SSRs and a corresponding lower abundance of SSRs per Mb of sequence than that of Garnica et al [27]. Preliminary analyses revealed a very limited number of loci containing SSRs that were common to the three *Phytophthora* species tested. The majority of the identified loci (81.6%) were common to *P. sojae* and *P. ramorum* only which is consistent with their closer phylogenetic relationship in clades 7 and 8 than to *P. infestans* in clade 1 [9,2,3]. Similarly, Garnica et al. found in their *in silico* analysis that 7.5% of their primers were, in theory, transferable between at least two species (mainly *P. ramorum* and *P. sojae*) and only 1.0% transferable between the three spe-

cies [27]. Among the selected sequences satisfying the above conditions, the number of repeats ranged from 3 to 17 and most SSRs showed seven repeats or less, with a repeat number of four being the most common in all species. The abundance of different repeat motifs differed slightly between species however, on average, (AAG)_n, (AGG)_n and (AGC)_n were the most abundant triplets in all three *Phytophthoras* (Fig. 5). These results differ from those reported by Garnica et al. in which (AGC)_n, (ACG)_n and (AGG)_n were the most abundant triplets amongst all the screened EST sequences [27]. It should, however be considered that unlike the study of Garnica our data are confined to SSR sequences for which it was possible to identify a homologue in at least one of the other two species. Therefore it could be hypothesised that motifs (AAG)_n and (AGG)_n are more abundant in more conserved genes. The dominance of trinucleotide SSRs compared to dinucleotide SSRs was not surprising considering that trinucleotides are abundant in coding regions of all higher eukaryotic genomes [31-33]. Dinucleotide repeats, in contrast, are characterised by higher mutation rates which may explain their abundance in introns and non-coding regions and lower frequency in coding regions, which cannot tolerate frame-shift mutations [34,35].

Table 7: Set of primers designed with the 2nd approach (Fig. 1B) to amplify genomic regions potentially containing SSRs in *Phytophthora* species of clade 7 [2].

Forward primer	Reverse Primer	SSRs	Source ^(a)	Gene ^(b)
S38F TCGTTTCTACGTGCTGGAY	S39R GTAGCACGCGAACATGAASA	(AAC) ₁₈	scaffold_26: 667079–667377	E
S40F TTCCTTAAGTGGGGGAGGAT	S41R TRTCGGCRITTCAGCTTCTGT	(AAG) ₄ ; (AAG) ₂₂	scaffold_125: 153837–154222	I
S42F GCTGCAAGAGTCSCTCGAGTA	S43R CTTGAGGATGTCRATGAGCA	(AG) ₂₁	scaffold_89: 132819–133248	I
S44F GTRGCTCCTTCCTTAAGTGG	S45R GTGCTGCASGTAYGGCTTC	(AAG) ₁₇	scaffold_75: 344500–344901	I
S46F GTTGCGCGTGAGGTTCTC	S47R CAAAGCTCTGCGTCC	(AG) ₂₂	scaffold_67: 282390–282656	I
S48F YCGGGCSACGGTAGG	S49R AAGAGCGTRAGCAGGAACC	(AG) ₁₈	scaffold_65: 225236–225440	I
S50F GTGGCTTCCACTGYTGCTG	S51R YATCAAGGACGTCAACTCGA	(AAC) ₉ ; (AACAGC) ₂₃	scaffold_48: 118994–119610	E
S52F CGGGATTTTRTCRGATCAGG	S53R CTGTYTGATCARCTCTCCGCT	(AGG) ₁₉	scaffold_46: 153312–153646	E
S56F CACGAGCTGCAGKCATAYCT	S57R AGAATKGAMGCGATCGAC	(AGG) ₁₆	scaffold_21: 370731–371140	E
S58F TCGATCRACAGAAGCTGCWA	S59R GGAGTTCGCCATCAACAAC	(AAG) ₁₄	scaffold_19: 606139–606624	I
S60F GGCGTTTAAAGCGTTTTAAA	S61R CGTCTTCTTCTTGACGCACA	(AC) ₁₈	scaffold_52: 422559–422915	I
S64F YTTGCGACTAGCAAAGTGG	S65R CGAACTCCTTGACAGGATGG	(AG) ₁₄	scaffold_56: 179585–179895	E
S66F GCAGYAGCCCCGGCCT	S67R GGAGTTCGCCATCAACAAC	(AAG) ₁₁	scaffold_12: 130593–130967	E
S68F CGTCGGTGGAGTAAACATCA	S69R AAAGGCGTTCGGAGAGYTG	(AG) ₁₄	scaffold_66: 83968–84423	I
S70F ATGACGAGGCAGCAGTTGAC	S71R AAGAACWGCSTACCTGCG	(ATC) ₁₃	scaffold_2: 564977–565285	I
S72F GCARCAATCTTCTGCTTYTTC	S73R ACACCTSCGTACWTTTCGTCA	(AAC) ₁₂	scaffold_92: 221631–221935	I
S74F CGGTGGTACTTGTCGTCCTC	S75R TSTCCGGCTACATCATCATC	(ATT) ₁₂	scaffold_41: 327977–328190	I
S76F GCATCTACGACCAGATCTACCC	S77R GTAGACSGAGATGATGGCGT	(AC) ₈	scaffold_127: 112530–112930	I

(a) = Gene sequences available at <http://genome.jgi-psf.org/sojae1/sojae1.home.html>.

(b) E = exons; I = Introns

Primers designed in the present study with the first approach were tested against a panel of 16 different *Phytophthora* species representing the breadth of diversity across the genus to amplify *P. sojae*, *P. ramorum* and *P. infestans* target regions containing 33, 17 and 8 SSRs respectively. Overall, these primers enabled the sequencing of 171 target regions which contained 211 SSRs ranging in repeat number from 3 to 16. Most of these SSRs showed seven repeats or less with four the most common repeat number and (AAG)_n, (AGG)_n and (AGC)_n the most common motifs. Trinucleotide repeats were dominant followed by pentanucleotide, tetranucleotide and dinucleotide repeats. This data indicate that such an approach can be useful to identify cross-specific SSR loci in the genus *Phytophthora*. As further genome sequences become available, for example, *P. capsici* <http://www.jgi.doe.gov/sequencing/why/CSP2006/Pcap>

[sici.html](#), the process can be refined to specific subsets of the genus. The mutation rates and, consequently, the practical utility of the identified SSRs in the study of the specific *Phytophthora* species need to be examined further. Undoubtedly, a risk of this approach is that the selection is biased towards more conserved sequences which may subsequently have a lower mutation rate that reduces their utility as polymorphic markers. Furthermore, the fact that *P. infestans* SSRs were all identified using primers designed on its own genome (Fig. 3) may indicate that this approach is less appropriate for distant relatives considering that, as stated above, *P. infestans* is phylogenetically distant from *P. sojae* and *P. ramorum*. However, the identification of intraspecific polymorphisms in some selected SSRs is encouraging and demonstrates that at least some of the selected SSRs are valuable for immediate practical applications (Table 6). This is consistent with the

Table 8: Accession numbers and SSRs for GenBank deposited sequences amplified using primers designed with the second approach (Fig. 1B).

Selected primers ^(a)	Phytophthora species				
	<i>P. alni</i> subsp. <i>alni</i> SCR2P2	<i>P. cambivora</i> SCR67	<i>P. europaea</i> SCR622	<i>P. fragariae</i> SCR333	<i>P. sojae</i> SCR555
S38-39	EF382833 No SSR	EF382832 No SSR	EF382831 No SSR	EF382830 No SSR	EF382829 (aac) ₁₈
S40-41	NS ⁺⁺	EF382801 (aac) ₄	EF382800 No SSR	EF382799 (aac) ₄	NS ⁺⁺
S42-43	NS ⁺⁺	EF382798 (ACG) ₆	EF382797 (acg) ₆ ; (agg) ₅	EF382796 No SSR	EF382795 (ag) ₂₁
S44-45	EF382793 (aac) ₄	EF382792 (aac) ₄	EF382794 No SSR	NS ⁺⁺	EF382791 (aag) ₅ ; (aag) ₅
S50-51	EF382790 Any SSR	NA ⁺	EF382789 No SSR	EF382788 No SSR	NS ⁺⁺
S52-53	NA ⁺	EF382786 No SSR	EF382787 No SSR	NS ⁺⁺	EF382785 (agg) ₁₉
S58-59	EF382784 (aac) ₄	EF382783 (aac) ₄	EF382782 No SSR	EF382781 (aac) ₄	EF382780 (aag) ₅ ; (aag) ₅
S64-65	EF382828 No SSR	EF382827 No SSR	EF382826 No SSR	EF382825 No SSR	EF382824 (ag) ₁₄
S68-69	EF382820 (aagg) ₄	EF382821 No SSR	EF382822 No SSR	EF382831 (aac) ₄ ; (aagg) ₄	EF382819 (ag) ₁₄
S70-71	EF382815 No SSR	EF382816 No SSR	EF382817 No SSR	EF382818 No SSR	EF382814 (act) ₁₃
S72-73	EF382810 No SSR	EF382811 No SSR	EF382812 No SSR	EF382813 No SSR	EF382809 (aac) ₁₂
S74-75	NS ⁺⁺	NS ⁺⁺	EF382807 No SSR	EF382808 No SSR	EF382806 (aat) ₁₂
S76-77	EF382802 No SSR	EF382803 No SSR	NS ⁺⁺	EF382805 No SSR	EF382804 (ac) ₈

^(a) = Primers listed in Table 7 and not reported in this table are those that did not produce any reliable sequence.
 NA⁺ = Isolate-primer combinations that did not produce any amplification or produced complex profiles (two or more fragments).
 NS⁺⁺ = Isolate-primer combinations for which single PCR bands were obtained but direct sequencing did not produced reliable results.

reported applicability of EST-SSRs across closely related taxa in other organisms as well as *Phytophthora* [23,36-38]. In the present study, the focus on the breadth of species (16) prevented the analyses of a wider number of target regions. However, the same method could be easily applied to the study of more regions from one or a few species.

The application of the first method enabled the identification of novel SSRs from all the 16 target species with those of clade 7 and 8 more highly represented (Fig. 3). A higher proportion of SSRs from species of the clade 7 and 8 was expected considering that *P. sojae* and *P. ramorum* belong to these two clades [2,3]. In light of this fact, a second approach to identify a greater number of polymorphic SSRs from within a more limited range of clade 7 taxa more closely related to *P. sojae* was investigated. Sixty *P. sojae* SSR candidates were compared by BLAST analysis against the complete genome sequence of the other two species yielding 18 SSR candidates which could be aligned

with homologous regions in *P. ramorum*. However in none of these 18 candidates (6 exons and 12 introns) was the SSR maintained in *P. ramorum*. In four of the more closely related species (*P. alni*, *P. cambivora*, *P. europaea* and *P. fragariae*), however, some of the SSR regions were conserved (Table 8). In this study the focus was on discovery of SSRs in invasive forest *Phytophthora* species within the clade 7a, perhaps a higher success rate in marker discovery would have followed a search amongst the closest related species in clade 7b (*P. sinensis*, *P. melonis*, *P. cajanae* and *P. vignae*) [2]. Although a few SSR markers with potential were discovered using this approach, it was not a highly efficient means of identifying new polymorphic SSR loci and highlights the lack of conservation of SSR loci, even amongst coding regions within a single ITS clade of *Phytophthora*. Some degree of cross-species amplification has been observed between SSRs in *P. infestans* with other Clade 1c taxa and it is therefore likely that a wider application of this method concentrated on the closest relatives would be more productive [23].

Conclusion

The present study has tested two different methods to generate SSR markers that can be utilised across a broad range of *Phytophthora* species. The final number of identified loci for any single species may not be sufficient to run a complete population genetics analysis and key studies on the inter- and intraspecific variation remain. A comprehensive dataset of candidate SSRs from a range of species has been created (Table 3, 4, 5). The detailed groundwork needed to amplify these regions from such a diverse collection of species and target regions has been completed which moves beyond the previous *in silico* approach to improve our understanding of the range and sequence conservation of SSR loci amongst species [27]. In general, the level of interspecific SSR sequence conservation, even amongst more closely related species within a single clade, was low and the method may not be the most efficient means of identifying novel SSR loci. Apart from their application as molecular markers, determining the abundance and density of SSRs in Oomycetes may help understand whether these sequences have any functional and evolutionary significance [17]. Furthermore, irrespective of the microsatellites, some of the amplified regions represent valuable marker regions for a number of applications [39]. A single optimal target gene for all *Phytophthora* species and assay requirements is unlikely to exist, therefore the continued identification and characterization of new target genes offers new opportunities for detection and phylogenetic studies [3,40,41].

Methods

Phytophthora isolates and DNA extractions

Twenty-eight isolates (16 *Phytophthora* species) sourced from the SCRI culture collection were used in this study (Table 1). Isolates and species were selected to represent taxa most relevant to European forestry that also represented the breadth of *Phytophthora* diversity defined according to clades based on ITS sequence analysis [2]. Isolates were stored on oatmeal agar at 5 °C and grown on French bean agar for routine stock cultures.

Total DNA was extracted from pure cultures of *Phytophthora* according to Schena and Cooke, diluted to 10 ng/μl and maintained at 5 °C for routine amplifications and at -20 °C for long term storage [42].

Analysis of sequences from *P. infestans*, *P. sojae* and *P. ramorum* genome projects and scanning for homologous SSRs

The predicted protein datasets of *P. infestans* (from the NCGR XGI database that was available prior to the Broad genome sequencing project) and *P. ramorum* and *P. sojae* <http://genome.jgi-psf.org/> were screened for SSR loci using Sputnik (Chris Abaijan <http://espressoftware.com/pages/sputnik.jsp>). Pairwise BLAST analysis

using the default parameters was used to select loci conserved in different species combinations [43]. Manual screening of these loci on the basis of SSR and flanking region DNA sequence conservation yielded a short-list for further analysis.

Primer design and amplification conditions

All primers (Table 2 and 7) were designed with the Primer3 Software set up to generate a T_m of 60 °C ± 2, a GC% between 20 and 80% and a length of 18–26 bp [44]. Primers were purchased from Eurogentec Ltd. (Belgium). Considerable effort was made to obtain successful amplification of single PCR bands from as many species as possible. This involved adjustment of MgCl₂ concentration (0.7, 1.0 or 1.7 mM) and annealing temperatures (55 or 58 °C) for PCR reactions (Table 3, 4, 5). Furthermore in some circumstances alternative primers were designed and tested to amplify the target regions from as many taxa as possible (Table 2). PCR reactions were performed in a total volume of 15 μl containing 10 ng of genomic DNA, 1.5 μl of 10× Reaction Buffer (Promega Corporation, WI, USA), 100 μM dNTPs, 0.7, 1 or 1.7 mM MgCl₂, 15 μg BSA, 2 unit of *Taq* polymerase (*Taq* DNA polymerase, Promega Corporation) and 1 μM of primers. PCR amplification conditions consisted of: 1 cycle of 95 °C for 2 min; 40 cycles of 94 °C for 30 s, 55 or 58 °C for 30 s, 72 °C for 60 s; and a final cycle of 72 °C for 5 min.

DNA sequencing

The best primers and amplification conditions were identified for all primer-species combinations and target DNA was re-amplified in a total volume of 50 μl to provide sufficient amplicon for direct sequencing. Single PCR bands were purified with the MinElute PCR Purification Kit (Qiagen Ltd. West Sussex, UK) to remove excess primers and nucleotides. Sequencing was carried out with the same primers utilized for the amplification in a dye-terminator cycle-sequencing reaction (FS sequencing kit, Applied Biosystems, Warrington, UK) and run on an ABI373 automated sequencer (Applied Biosystems). All selected PCR fragments were sequenced using both the forward and the reverse primers.

Sequence analysis and SSRs scanning

The "Sequence Navigator" software (Applied Biosystems) was utilised to evaluate reliability of sequences and to compare forward and reverse sequences to create a consensus sequence. Non-reliable sequences in which both forward and reverse sequences contained doubtful bases were discarded. All sequences obtained in the present study were also parsed to a web version of SPUTNIK <http://cbl.jabri.fr/outils/Pise/sputnik.html>, which uses a recursive algorithm to search for repeated patterns of nucleotides of length between 2 and 5.

Authors' contributions

LS conceived the study, carried out the molecular analyses, optimized the described method, performed data acquisition, data analysis and data interpretation and wrote the manuscript.

LM performed analysis of sequences from genome projects and scanned for homologous SSRs.

DEL coordinated, supervised and contributed to the design of the study, provided intellectual input to optimize the method and revised the manuscript critically.

Acknowledgements

This work was funded by the European Union with a Marie Curie Intra-European Fellowship (Project: MEIFCT-2003-502327) and by the Scottish Government Rural and Environment Research and Analysis Directorate (RERAD). We thank C.M. Brasier and others for providing isolates, N. Williams for invaluable help with the culture collection and Nik Grunwald for providing a list of *P. sojae* SSR markers. Cultures are held under SASA License No. PH/11/2008.

References

- Baldauf SL, Roger AJ, Wenk-Siefert I, Doolittle WF: **A Kingdom-level phylogeny of eukaryotes based on combined protein data.** *Science* 2000, **290**:972-977.
- Cooke DEL, Drenth A, Duncan JM, Wagels G, Brasier CM: **A molecular phylogeny of *Phytophthora* and related Oomycetes.** *Fungal Genet Biol* 2000, **30**:17-32.
- Kroon LPNM, Bakker FT, Bosch GBM van den, Bonants PJM, Flier WG: **Phylogenetic analysis of *Phytophthora* species based on mitochondrial and nuclear DNA sequences.** *Fungal Genet Biol* 2004, **41**:766-782.
- Tyler BM, Tripathy S, Zhang X, Dehal P, Jiang RHY, Aerts A, Arredondo F, Baxter L, Damasceno CMB, Dickerman A, Dorrance AE, Dou D, Dubchak I, Garbelotto M, Gijzen M, Gordon S, Govers F, Grunwald N, Huang W, Ivors K, Jones RW, Kamoun S, Krampis K, Lamour K, Lee MK, McDonald WH, Medina M, Meijer HJG, Nordberg E, MacLean DJ, Ospina-Giraldo MD, Morris P, Phuntumart V, Putnam N, Rash S, Rose JKC, Sakihama Y, Salamov AA, Savidor A, Scheuring C, Smith B, Sobral BWS, Terry A, Torto-Alalibo T, Win J, Xu Z, Zhang H, Grigoriev I, Rokhsar D, Boore J: ***Phytophthora* genome sequences uncover evolutionary origins and mechanisms of pathogenesis.** *Science* 2006, **313**:1261-1265.
- Erwin DC, Ribeiro OK: ***Phytophthora* diseases worldwide.** APS Press, St Paul, MN, USA; 1996.
- Jurskis V: **Eucalypt decline in Australia, and a general concept of tree decline and dieback.** *Forest Ecol Manag* 2005, **215**:1-20.
- Ristaino JB, Gumpertz ML: **New frontiers in the study of dispersal and spatial analysis of epidemics caused by species in the genus *Phytophthora*.** *Annu Rev Phytopathol* 2000, **38**:541-576.
- Rizzo DM, Garbelotto M, Hansen EM: ***Phytophthora ramorum*: integrative research and management of an emerging pathogen in California and Oregon forests.** *Annu Rev Phytopathol* 2005, **43**:309-335.
- Blair JE, Coffey MD, Park S, Geiser DM, Kang S: **A multi-locus phylogeny for *Phytophthora* utilizing markers derived from complete genome sequences.** *Fungal Genet Biol* 2008, **45**:266-277.
- De Merlier D, Chandelier A, Debruxelles N, Noldus M, Laurent F, Dufays E, Claessens H, Cavelier M: **Characterization of alder *Phytophthora* isolates from Wallonia and development of SCAR primers for their specific detection.** *J Phytopathol* 2005, **153**:99-107.
- Cooke DEL, Young V, Birch PRJ, Toth R, Gourlay F, Day JP, Carnegie S, Duncan JM: **Phenotypic and genotypic diversity of *Phytophthora infestans* populations in Scotland (1995-97).** *Plant Pathol* 2003, **52**:181-92.
- Ivors KL, Hayden K, Bonants PJM, Rizzo DM, Garbelotto M: **AFLP and phylogenetic analyses of North American and European populations of *Phytophthora ramorum*.** *Mycol Res* 2004, **108**:378-392.
- Cooke DEL, Lees AK: **Markers, old and new, for examining *Phytophthora infestans* diversity.** *Plant Pathol* 2004, **53**:692-704.
- Selkoe KA, Toonen RJ: **Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers.** *Ecol Lett* 2006, **9**:615-629.
- Mascheretti S, Croucher PJP, Vettraino A, Prospero S, Garbelotto M: **Reconstruction of the Sudden Oak Death epidemic in California through microsatellite analysis of the pathogen *Phytophthora ramorum*.** *Mol Ecol* 2008, **17**:2755-2768.
- Li YC, Korol AB, Fahima T, Beiles A, Nevo E: **Microsatellites: genomic distribution, putative functions, and mutational mechanisms: A review.** *Mol Ecol* 2002, **11**:2453-2465.
- Toth G, Gaspari Z, Jurka J: **Microsatellites in different eukaryotic genomes: survey and analysis.** *Genome Res* 2000, **10**:967-981.
- Chistiakov DA, Hellems B, Volckaert FAM: **Microsatellites and their genomic distribution, evolution, function and applications: A review with special reference to fish genetics.** *Aquaculture* 2006, **255**:1-29.
- Gobbin D, Pertot I, Gessler C: **Identification of microsatellite markers for *Plasmopara viticola* and establishment of high-throughput method for SSR analysis.** *Eur J Plant Pathol* 2003, **109**:153-164.
- Dobrowolski MP, Tommerup IC, Blakeman HD, O'Brien PA: **Non-mendelian inheritance revealed in a genetic analysis of sexual progeny of *Phytophthora cinnamomi* with microsatellite markers.** *Fungal Genet Biol* 2002, **35**:197-212.
- Dobrowolski MP, Tommerup IC, Shearer BL, O'Brien PA: **Three clonal lineages of *Phytophthora cinnamomi* in Australia revealed by microsatellites.** *Phytopathology* 2003, **93**:695-704.
- Knapova G, Gisi U: **Phenotypic and genotypic structure of *Phytophthora infestans* populations on potato and tomato in France and Switzerland.** *Plant Pathol* 2002, **51**:641-653.
- Lees AK, Wattier R, Shaw DS, Sullivan L, Williams NA, Cooke DEL: **Novel microsatellite markers for the analysis of *Phytophthora infestans* populations.** *Plant Pathol* 2006, **55**:311-319.
- Ivors K, Garbelotto M, Vries ID, Ruyter-Spira C, Te Hekker B, Rosenzweig N, Bonants P: **Microsatellite markers identify three lineages of *Phytophthora ramorum* in US nurseries, yet single lineages in US forest and European nursery populations.** *Mol Ecol* 2006, **15**:1493-505.
- Prospero S, Black JA, Winton ML: **Isolation and characterization of microsatellite markers in *Phytophthora ramorum*, the causal agent of sudden oak death.** *Mol Ecol* 2004, **4**:672-674.
- Glenn TC, Schable NA: **Isolating microsatellite DNA loci.** In *Molecular evolution: producing the biochemical data, part B* Edited by: Zimmer EA, Roalson E. Academic Press, San Diego, USA; 2005:202-222.
- Garnica DP, Pinzón AM, Quesada-Ocampo LM, Bernal AJ, Barreto E, Grünwald NJ, Restrepo S: **Survey and analysis of microsatellites from transcript sequences in *Phytophthora* species: frequency, distribution, and potential as markers for the genus.** *BMC Genomics* 2006, **7**:245.
- Robinson AJ, Love CG, Batley J, Barker G, Edwards D: **Simple sequence repeat marker loci discovery using SSR primer.** *Bioinformatics* 2004, **20**:1475-1476.
- Prospero S, Hansen EM, Grünwald NJ, Winton LM: **Population dynamics of the sudden oak death pathogen *Phytophthora ramorum* in Oregon from 2001 to 2004.** *Mol Ecol* 2007, **16**:2958-2973.
- Widmark AK, Andersson B, Cassel-Lundhagen A, Sandström M, Yuen JE: ***Phytophthora infestans* in a single field in southwest Sweden early in spring: symptoms, spatial distribution and genotypic variation.** *Plant Pathol* 2007, **56**:573-579.
- Borstnik B, Pumpernik D: **Tandem repeats in protein coding regions of primate genes.** *Genome Res* 2002, **12**:909-915.
- Metzgar D, Bytof J, Wills C: **Selection against frameshift mutations limits microsatellite expansion in coding DNA.** *Genome Res* 2000, **10**:72-80.
- Subramanian S, Madgula VM, George R, Mishra RK, Pandit MW, Kumar CS, Singh L: **Triplet repeats in human genome: distribution and their association with genes and other genomic regions.** *Bioinformatics* 2003, **19**:549-552.

34. Karaoglu H, Lee CMY, Meyer W: **Survey of simple sequence repeats in completed fungal genomes.** *Mol Biol Evol* 2004, **22**:639-649.
35. Katti MV, Ranjekar PK, Gupta VS: **Differential distribution of simple sequence repeats in eukaryotic genome sequences.** *Mol Biol Evol* 2001, **18**:1161-1167.
36. Cordeiro GM, Casu R, McIntyre CL, Manners JM, Henry RJ: **Microsatellite markers from sugarcane (*Saccharum* spp.) ESTs cross transferable to *Erianthus* and *Sorghum*.** *Plant Sci* 2001, **160**:1115-1123.
37. Woodhead M, Russell J, Squirrel J, Hollingsworth PM, Cardle L, Ramsay L, Gibby M, Powell W: **Development of EST-SSRs from the Alpine lady-fern, *Athyrium distentifolium*.** *Mol Ecol Notes* 2003, **3**:287-290.
38. Zhan A, Bao ZM, Wang XL, Hu JJ: **Microsatellite markers derived from bay scallop *Argopecten irradians* expressed sequence tags.** *Fisheries Science* 2005, **71**:1341-1346.
39. Cooke DEL, Schena L, Cacciola SO: **Tools to detect, identify and monitor *Phytophthora* species in natural ecosystems.** *J Plant Pathol* 2007, **89**:13-28.
40. Schena L, Hughes KJD, Cooke DEL: **Detection and quantification of *P. ramorum*, *P. kernoviae*, *P. citricola* and *P. quercina* in symptomatic leaves by multiplex real-time PCR.** *Mol Plant Pathol* 2006, **7**:365-379.
41. Schena L, Duncan JM, Cooke DEL: **Development and application of a PCR based "molecular tool box" for the detection of *Phytophthora* species damaging forests and natural ecosystems.** *Plant Pathol* 2008, **57**:64-75.
42. Schena L, Cooke DEL: **Assessing the potential of regions of the nuclear and mitochondrial genome to develop a "molecular tool box" for the detection and characterization of *Phytophthora* species.** *J Microbiol Methods* 2006, **67**(1):70-85.
43. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215**:403-410.
44. Rozen S, Skaletsky HJ: **Primer3 on the WWW for general users and for biologist programmers.** *Methods Mol Biol* 2000, **132**:365-386.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

