



Characterization of *Phytophthora* de Bary inciting capsule rot/azhukal and leaf blight disease of cardamom (*Elettaria cardamomum* Maton)

*Susheela Bhai R and Sarma Y R

*Indian Institute of Spices Research, Marikunnu.P.O, Calicut - 673 012, Kerala, India.

(Manuscript Received : 18-03-2005; Revised : 23-11-2005; Accepted : 28-12-2005)

Abstract

A detailed study was carried out to characterize the *Phytophthora* species associated with capsule rot and leaf blight diseases in cardamom. *Phytophthora* was isolated from rot affected plant parts such as leaf, capsule and pseudostem collected from different cardamom growing tracts of Idukki and Wyanad districts of Kerala during the monsoon and post monsoon periods. Morphological and physiological studies such as colony morphology, growth rate, morphology of sporangia, pedicel length, caducity, chlamydospores, sex organs, compatibility type, growth on synthetic medium and temperature relationship was studied using 25 *Phytophthora* isolates from cardamom. Two groups of *Phytophthora* could be identified from infected capsules, leaves and pseudo stem. The group one having pigmentation, chlamydospore production and growth at >35 °C where as the group two having no pigmentation, no chlamydospore production and no growth at 35 °C. The group one is identified as *P. nicotianae* var. *nicotianae* and group two as *P. meadii*. From the studies it has been confirmed that two species of *Phytophthora* viz. *P. meadii*, and *P. nicotianae* var. *nicotianae* are involved in initiating capsule rot occurring during the monsoon season of which *P. meadii* is the predominant one, The leaf blight occurring during the post monsoon season is due to *P. nicotianae* var. *nicotianae*. However, there is certain amount of overlapping of these in initiating the disease in cardamom. *P. palmivora* could not be isolated from the infected plant parts as reported earlier.

Key words: Azhukal, characterization, *Elettaria cardamomum*, *Phytophthora meadii*, pigmentation, sporangia

Introduction

Capsule rot (Azhukal) disease caused by *Phytophthora* species has been implicated as a serious problem causing extensive damage in many cardamom growing tracts in the high ranges of Kerala. The disease appears with the onset of the southwest monsoon season. It is characterized by the rotting of leaves, panicles, capsules and pseudostem. Menon *et al.*, (1972) for the first time reported the disease as caused by *Phytophthora* and proved Koch's postulates and identified the pathogen as *P. nicotianae* (Thankamma and Pillai, 1973; Liyanage *et al.*, 1983) *P. palmivora* (Radha and Joseph, 1974) and *P. meadii* (Sastry, 1982., Anon 1986). Later another disease called leaf blight due to *P. meadii* was also identified (Anon 1986). This disease appears during the post monsoon period from November to February.

The involvement of various species of *Phytophthora* in causing capsule rot and leaf blight disease as reported by various workers is controversial and an elaborate study in this aspect needs utmost importance. The present study was taken up with the objective to identify the actual pathogen involved in initiating capsule rot and leaf blight disease of cardamom and the results are discussed.

Materials and Methods

Azhukal/ rot affected plant parts such as capsules, pseudostem and leaves were collected from different cardamom growing tracts during the monsoon season. Leaf blight samples were collected during the post monsoon season. The isolation of the pathogen was done using PVPH medium (Tsao and Guy, 1977). Around 120 isolates were collected from 80 different locations of which 25 isolates were short listed and studied in detail

for colony morphology, growth rate on agar, sporangial morphology, ontogeny, caducity, chlamyospore production, sex organs formation, pigmentation and temperature relations (Table 1)

in dark at $24 \pm 1^\circ\text{C}$ and the growth marked at 24hrs Interval for 120 hrs. After the incubation period two measurements of the colony diameter were made at right angles to each other and the growth rate was calculated and expressed as mm day^{-1}

Table 1. Morphology of *Phytophthora* isolates of cardamom

| Place of Collection | Plant part | Isolate No. | Colony morphology/Pattern |
|---------------------|------------|-------------|--|
| Myladumpara | Capsule | CPC 159 | Chrysanthemum |
| Pachakkanam | Pseudostem | CPP 244 | Uniform cottonwool like aerial mycelium with slight lobes |
| Pallivasal | Capsule | CPC 175 | Uniform cottonwool like aerial mycelium with slight lobes |
| Udumbanchola | Capsule | CPC 109 | Uniform cottonwool like aerial mycelium with slight lobes |
| Mali | Capsule | CPC 310 | Lobed |
| Puttady | Capsule | CPC 311 | Chrysanthemum |
| Vandanmedu | Leaf | CPL 312(1) | Uniform cotton wool like aerial mycelium with slight lobes |
| Vandanmedu | Pseudostem | CPP 312(d) | Lobed |
| Pulyanmala | Capsule | CPC 313 | Lobed |
| Mettukuzhy | Capsule | CPC 317 | Uniform cotton wool like aerial mycelium with slight lobes |
| Myladumpara | Capsule | CPC 318 | Lobed |
| Poopara | Leaf | CPI.342 | Uniform cotton wool like aerial mycelium with slight lobes |
| Puttady | Capsule | CPC 349 | Lobed |
| Myladumpara | Pseudostem | CPP 350 | Uniform cotton wool like aerial mycelium with slight lobes |
| Myladumpara | Leaf | CPL 352 | Uniform cotton wool like aerial mycelium with slight lobes |
| Pallivasal | Capsule | CPC353 | Lobed |
| Muthukad | Pseudostem | CPP 356 | Lobed |
| Muthukad | Capsule | CPC 357 | Lobed |
| Udumbanchola | Capsule | CPC 358 | Lobed |
| Myladumpara | Capsule | CPC 363 | Chrysanthemum |
| Parathode | Capsule | CPC 365 | Lobed |
| Meppadi | Leaf | CPL 366 | Lobed |
| Meppadi | Leaf | CPL 368 | Chrysanthemum |
| Meppadi | Leaf | CPL 369 | Uniform cotton wool like aerial mycelium with slight lobes |
| Thachungal | Capsule | CPC 375 | Chrysanthemum |

Colony morphology and growth rate on agar

The colony morphology of all the 25 isolates were studied in carrot agar and corn meal agar (Brasier and Griffin, 1979; Liyanage and Wheeler, 1989). Four mm discs cut from the edge of 72 h old culture (of all the 25 isolates) was inoculated centrally in 90 mm diameter plates containing 15ml of the corresponding medium, covered in aluminium foil and incubated in total darkness at $24 \pm 1^\circ\text{C}$ for 5 days. Colony morphology was examined against a black background.

The growth rate was studied using five different types of media viz., carrot agar (Brasier *et al.*, 1993), capsule extract agar, oat meal agar and casein hydrolysate tyrosine agar or Timmer's medium (Timmer *et al.*, 1970). Capsule extract agar was prepared by smashing 200 g tender capsules and boiling in 1 litre of distilled water for 10 min. The juice was squeezed through two layers of cheese cloth and volume made up to one liter. Agar was added @17g /l and sterilized. For studying the growth rate, isolates were inoculated centrally on plates (90mm diameter) containing 15ml of the medium and incubated

Sporangial morphology, ontogeny and measurements

Sporangial morphology and sporulation was studied using slide culture technique in carrot agar medium (Dhantanarayana *et al.*, 1984). Inoculum plugs of 3 mm size were cut from the edge of 72 h old culture and placed on the agar-coated slide and incubated for 5-7 days at $24 \pm 1^\circ\text{C}$ and then fixed in lactophenol. These slides were microscopically examined for sporangial morphology and ontogeny.

Sporangial measurements were taken as per Tsao *et al.* (1985). Discs of 4mm cut from the edge of 72hrs old actively growing culture incubated in the dark was placed in 5ml of sterile distilled water in petri-dishes and incubated under continuous light for five days. The sporangia bearing mycelium that had grown out into the surrounding water during the incubation period were cut with a scalpel and mounted on a drop of lactophenol. The length, breadth and pedicel length of 50 randomly chosen sporangia was measured under 15x X 40x magnification using an ocular micrometer.

Caducity

Among the discs incubated under light, one disc each was transferred into a test tube with 4ml of sterile water and shaken for 1 min to dislodge the sporangia. This water was poured into a counting dish and the total number of dislodged sporangia was counted under 4x magnification. The disc after dislodging the sporangia was pressed against a slide and the number of attached sporangia/microscopic field was counted. The total number of sporangia/disc was calculated and percent caducity was calculated using the formula

$$\frac{\text{No. of detached sporangia/disc}}{\text{Total no. of sporangia / disc}} \times 100$$

Chlamydospore production

Ribeiro's liquid medium (Ribeiro *et al.*, 1975) @ of 50ml/ 100ml flask was sterilized at 120°C for 20 min. These were inoculated with 5mm discs of 72 h old culture and incubated in the dark at 24±1°C for one month. Similarly 5mm discs were inoculated in carrot agar and Ribeiro's solid agar medium and incubated in the dark for one month at 24±1°C. Infected plant parts were also stored in the refrigerator at 10°C and observed one month after incubation

Sexual reproduction and mating type

Dual culture method was used for sex organ production. Test isolates of cardamom were plated against known A1 isolate of black pepper in carrot agar with thiamine (Brasier *et al.*, 1993) and also in Ribeiro's medium. For pairing, 3mm discs cut from the edge of 72 h old culture of both A1 and test isolates of cardamom were placed on opposite sides of the petri dish at a distance of 3cm apart. The plates were incubated in the dark at 21±1°C. Observations were taken at 5 day intervals for 21 days. Measurements of oogonia and antheridia were also taken from these paired plate cultures.

Pigmentation

Timmers's medium containing casein hydrolysate and tyrosine were used for pigment production. Agar slopes of the medium were made and inoculated with 3 mm discs of 72 h old culture and incubated in the dark at 24±1°C for 15-21 days (Kennedy and Duncan, 1995). Pigment production was observed on the 21st day

Temperature relationship

Inoculum discs of 3mm were cut from the margin of 72 h old culture and inoculated centrally in 90 mm petri dishes containing 15 ml of the medium. These plates were

incubated in a BOD incubator at temperatures ranging from 10-35°C and observations on the radial growth was measured at 24 hr interval for 96 hr and the average growth rate was calculated.

Results

Colony morphology and growth rate

Three distinct type of colony morphology was mainly observed among the isolates. They were uniform cotton wool like aerial mycelium over the entire colony, stellate or petalloid pattern and lobed pattern (Plate 1). No particular pattern could be assigned to capsule, leaf or pseudo stem isolates.




Plate 1. Colony morphology of *Phytophthora* isolates of cardamom

Among the five media tested viz. carrot agar, corn meal agar, capsule extract agar, casein hydrolysate tyrosine agar and oat meal agar, carrot agar was found to be the best medium for the growth (7-9 mm/d) followed by oat meal agar. In casein hydrolysate tyrosine agar and capsule extract agar, the growth rates of the isolates were very poor (1.5 mm/d). Pseudostem isolates showed comparatively faster growth (4.35 mm/d) followed by leaf isolates (3.9mm/d) and capsule isolates (Table 2).

Sporangial morphology and ontogeny

Sporangia, irrespective of whether capsule, pseudostem or leaf isolate, were ovoid to ellipsoidal with basal plug conspicuous in some. The papillae were very

prominent with wide exit pores. Sporangia were mostly symmetrical. Although asymmetrical types were also noticed occasionally. In some isolates swelling at the point of sporangial formation was observed. Sporangia with round base and some times with slight tapering was also noticed. Sporangia were produced in simple sympodia and well defined in agar. They were formed in abundance when the culture disc was kept in water under continuous fluorescent light (Plate 2). Certain isolates showed the presence of hyphal thickenings and sporangial outgrowths (Plate 2a & 2b).




Plate 2a. Sporangial morphology of *Phytophthora* isolates of cardamom

Sporangial measurements:-

Length, breadth and l/b ratio

The size of the sporangia differs considerably in solid agar and in water culture. Sporangia were generally larger in solid culture. There was no considerable variation in sporangial size between capsule, leaf and pseudostem isolates. The mean sporangial length: breadth ratio of

Table 2. Growth rate on Agar media

| Media and time after inoculation | Capsule | | Leaf | | Pseudo-stem | |
|----------------------------------|------------|-------------|------------|-------------|-------------|-------------|
| | Mean | Range | Mean | Range | Mean | Range |
| CA 0-120h | 7.92±0.22 | (6.7-9.00) | 8.06±.41 | (6.75-9.00) | 8.6±0.25 | (7.9-9.00) |
| CMA 0-120h | 5.54±0.24 | (4.54-6.18) | 5.92±0.18 | (5.43-6.43) | 6.04±0.20 | (5.86-6.43) |
| CHT 0-120h | 3.13± 1.54 | (2.15-4.97) | 3.90±0.50 | (3.86-4.03) | 4.35±0.25 | (4.00-5.08) |
| CEA 0-120h | 4.32±0.20 | (2.82-6.13) | 4.44±0.13 | (3.93-4.63) | 4.19±0.17 | (3.82-4.5) |
| OMA 0-120h | 7.29±0.23 | (5.69-7.69) | 7.32± 0.17 | (6.69-7.69) | 6.83±0.12 | (6.57-7.07) |




Plate 2b. Sporangial ontogeny of *Phytophthora* isolates of cardamom

the three group of isolates were almost same and were more than 1.7 (1.7-2.09). The sporangial dimensions of individual cultures in solid and water culture is given in Table 3.

The mean sporangial length in solid culture ranged from 51.39-56.82 μm and breadth ranged from 27.43-28.45 μm . In water culture it ranged from 42.22-45.40 μm and 24.99-25.09 μm respectively. The l/b ratio in solid culture ranged from 1.84-2.09 as compared to 1.75-1.82 in water (Table 3).

Pedicle length

In all the cases the pedicle length was found in the intermediate group range ie. 5-20 μm . But certain isolates showed considerable variation in pedicle length in solid and water culture (Table .3). The mean pedicle length of

Table 3. Mean sporangial dimensions of *Phytophthora* isolates of cardamom

| Variate | Capsule | | Leaf | | Pseudostem | |
|----------------|---------------|----------------|---------------|----------------|---------------|----------------|
| | Solid culture | Liquid culture | Solid culture | Liquid culture | Solid culture | Liquid culture |
| Length | 55.25±3.18 | 45.40±1.56 | 51.39±3.05 | 42.22±2.96 | 56.82±4.05 | 43.85±2.77 |
| Range | 25.8±113.52 | 20.64±116.1 | 33.54±70.2 | 25.8±72.24 | 29.7±108.36 | 25.8±79.98 |
| Breadth | 28.36±1.09 | 24.99±0.56 | 28.45±1.30 | 24.79± 1.84 | 27.43±0.74 | 25.0± 1.63 |
| Range | 16.2±46.44 | 10.32±41.28 | 18.06±43.86 | 51.16±41.28 | 154.48±36.12 | 15.48±38.7 |
| L/B | 1.95±0.09 | 1.82±0.04 | 1.84±0.10 | 1.75±0.09 | 2.09±0.16 | 1.77±0.53 |
| Range | 1.62-2.91 | 1.60-2.11 | 1.59-2.21 | 1.56-2.08 | 1.82-2.52 | 1.63-1.87 |
| Pedicle length | 15.05-1.40 | 21.32±1.45 | 15.40±3.75 | 21.44±0.66 | 14.83±2.09 | 23.96±3.31 |
| Range | 2.58-67.08 | 1.29-69.66 | 5.16-45.9 | 5.16-67.08 | 2.58-45.9 | 2.58-69.66 |

the three group of isolates in solid culture varied from 14.83-16.62 μm and in water culture it varied from 21.32-23.96 μm . The high frequency of mean pedicel length in the range of 5-20 μm was noticed in solid culture. The pedicel length of >20 μm was observed in the range of 4-68%. In water culture the pedicel length in the range of 5-20 μm was noticed @ 53.8% and >20 μm was noticed @ 46.2%. The capsule isolates showed a frequency of 50-96% and mean pedicel length in the range of 5-20 μm .

Caducity

The degree of caducity (expressed as % detached sporangia) varied in solid and water culture. Caducity was more in water culture (28.46-41.72%) as compared to solid culture (11.57-14.55%) (Table 4). Not much variation was observed in percent caducity between the three groups of isolates in solid culture. In water culture pseudostem isolates were found to be comparatively more caducous (41.72%) compared to capsule isolates (35.84%). Leaf isolates showed the least caducity (28.46%).

Among the capsule isolates CPC 159, CPC 309, CPC 311, CPC 313 and CPC 349 showed very low caducity (below 10%) in solid culture while others have an average caducity of 18.5%. In water culture these isolates showed very high caducity (more than 25%). But the isolate CPC 313 was found to have low caducity in water when compared to others (16.23%). In solid culture, the three groups of isolates showed almost uniform caducity (11.57-14.55%).

Chlamyospore production

Among the three sources, it was observed that capsule isolates were very poor in producing chlamyospores as compared to leaf and pseudostem isolates. Only 31.25% of the capsule isolates produced chlamyospores, whereas 60% of the leaf and 50% of the pseudostem isolates produced chlamyospores in culture. Some of the isolates produced chlamyospores in water. Both

terminal and intercalary chlamyospores were present. They were spherical, hyaline and thin walled and the diameter ranged from 20-31 μm with an average of 26.63 μm (Plate 3).




Plate-3 Chlamyospores of *Phytophthora* isolates of cardamom

Sexual Reproduction

All the isolates except two produced sex organs on pairing with A1 isolate of *P. capsici* of black pepper. Certain isolates produced sex organs on a band formed in the junction of meeting the two isolates. Oogonia were spherical, hyaline and thin walled but in some isolates it became golden yellow in colour (Plate 4). Antheridia are purely amphigynous. Some of the isolates produced sex organs within 5-8 days whereas others took 15-21 days. Among the three sources it was observed that pseudostem isolates showed the biggest oogonia (33.23 μm) and the smallest antheridia (12.35 μm) whereas the leaf isolates showed the smallest oogonia and the biggest antheridia (17.13 μm). In capsule isolates the l/b of antheridia was almost equal to one (0.97) whereas it was

1.19 and 0.85 for leaf and pseudostem isolates respectively (Table 5). The oogonial diameter of capsule isolates ranged from 24.04-39.69 μm and the antheridial length ranged from 11.29 μm -16.77 μm . Of the 25 isolates examined, 24 belonged to the A2 compatibility type and one isolate was found sterile with out producing any oogonia or antheridia even in pairing (CPC 375). The mating behaviour was also variable with respect to isolates. Some of the isolates produced few oogonia while others produced oogonia in abundance.




Plate-5 Pigmentation in Timmers medium

Temperature relationship

All the isolates were found to be growing between 10-32°C whereas they failed to grow at 5°C or at 40°C. Most of the pseudo stem and leaf isolates grew well at 36°C. The optimum temperature range for growth of most of the isolates has been found to be between 20°C and 28°C. Irrespective of the source of isolation, maximum growth (8.9mm/day) of the isolates occurred at 20-28°C and grew up to 32°C. At 32°C growth was observed at the rate of 1-6mm/d and the capsule isolates showed the minimum growth (1.89mm/d) as compared to leaf and pseudostem isolates (4.54 mm/d). Minimum growth was observed at 10°C (1-1.73 mm/d) for all the three groups.

Table 5. Measurements of oogonia and antheridia of *Phytophthora* isolates cardamom

| Isolates | Antheridia (μm) | | Oogonia μm Diameter |
|------------|------------------------------|------------------|-----------------------------------|
| | Length | Breadth | |
| Capsule | 14.32 \pm 0.42 | 14.72 \pm 0.25 | 31.16 \pm 1.29 |
| Leaf | 17.13 \pm 3.48 | 14.37 \pm 1.08 | 29.94 \pm 1.18 |
| Pseudostem | 12.35 \pm 0.85 | 14.59 \pm 0.72 | 33.23 \pm 2.56 |

Discussion .

Successful isolations of pure cultures of *Phytophthora* from infected tissues of cardamom using PVPH and BNPR+HMI medium (Nair and Menon, 1980) and Lima Bean Agar medium (Thankamma *et al.*, 1968) was reported earlier. In cardamom three distinct types of colony morphology were observed among the isolates. They were uniform cotton wool like aerial mycelium over entire colony with vague lobed pattern, stellate or petalloid pattern and lobed pattern. The result is in accordance with those obtained for colonies of M isolates of rubber described by Dantanarayana *et al.* (1984) and also S & L types described by Brasier and Griffin (1979). The variability in colony morphology is found to be a distinctive character among the isolates of cardamom and it also varied with type of media used. No definite pattern was noticed specifically for capsule, leaf or pseudostem isolates. Incidentally the isolates exhibited variability in colony morphology and hence could not be used as an important taxonomic character.

resent.
nd the
26.63




Plate-4 Oogonia and oospores of *Phytophthora* isolates of cardamom

Table 4. Mean caducity and pedicel length of *Phytophthora* isolates of cardamom

| Isolates | Caducity% | | Pedicel length(μm) | |
|-------------|------------------|------------------|----------------------------------|------------------------------------|
| | Solid culture | Water culture | Solid culture | Water culture |
| Capsule | 14.55 \pm 1.82 | 35.34 \pm 3.13 | 16.62 \pm 1.06 (7.74-24.04) | 21.32 \pm 1.47 (11.54-81.825) |
| Leaf | 11.57 \pm 3.04 | 28.46 \pm 2.89 | 15.40 \pm 3.74 (8.31-24.95) | 21.43 \pm 0.66 (19.24-22.53) |
| Pseudo-stem | 13.49 \pm 4.23 | 41.72 \pm 4.62 | 14.83 \pm 2.33 (9.06-19.37) | 23.96 \pm 3.31 (17.33-32.8) |

Pigment production

Chocolate brown pigmentation was observed in Timmer's medium (CHT) by some of the isolates when incubated for 21 days at 24-28°C whereas others had no pigmentation in the medium. (Plate-5).

is on
pper.
med
were
tes it
a are
l sex
lays.
stem
of the
lates
gest
b of
was

In the present study the growth rate of isolates showed considerable variation in different media used. Variability in growth rate was reported among *P. meadii* and *P. palmivora* (M&P) isolates of rubber and *P. palmivora* and *P. megakarya* (M&P) isolates of Cocoa (Brasier and Griffin, 1979, Dantanarayana *et al.*, 1984). Variability in growth rate was also reported for isolates collected from various diseased parts of rubber (Liyange and Wheeler, 1989). Thus the variation in growth rate with respect to media, a condition under which it was grown and the parts from which isolations have been made was true in the case of cardamom isolates also.

The pedicel length of the isolates is in agreement with the key's of Waterhouse (1963), Newhook *et al* (1978) Stamps *et al* (1990), and Ho (1981). Waterhouse (1965) described *P. meadii* as having deciduous sporangia with pedicel 2-10 μm . Newhook *et al* (1978) and Stamps *et al.* (1990) described the pedicel as caducous small, medium or long. Cardamom isolates came under the second category of medium pedicel length. According to the key of Ho (1981) caducous sporangium having pedicel length of 5-20 μm which sheds in water was reported for species such as *P. botryos*, *P. colocasiae* Raciborshi, *P. illicis* Buddenhagen and Young, *P. meadii* Mc Rae and *P. megakarya* (Brasier and Griffin 1979). The mean pedicel length fell a between 15-24 μm which corresponded to the intermediate pedicel length group as described by Zentmyer *et al.*, (1977) and Al-Hedaithy and Tsao (1979a). The mean l/b ratio ranged from 1.77-2.09. The l/b ratio of cardamom isolates were >1.6 which included both *P. meadii* and *P. nicotianae* var. *nicotianae*. The l/b ratio of *P. nicotianae* as described by Garreston-Cornell (1989) was 1.3-1.5. All the isolates sporulate heavily in water in light whereas sporulation in the dark in both solid and water culture was very scanty. Only a few sporangia were produced in solid agar in dark except one isolate (CPC342) which produced plenty of sporangia in solid agar in dark with intercalary chlamydo spores and junctional thickenings. The present study was similar to those reported by Rajalekshmy and Annakutty Joseph (1986). They observed scanty sporangial production in plate cultures of *P. meadii* incubated in the dark.

The caducity in water (28.46-41.72%) was double than that in solid agar plate method (11.57-14.%) which corresponded with the increase in pedicel length. The results obtained in the present study is in contrast to that of *P. palmivora* where the percentage detached sporangia in solid agar plate method was higher than in agar disc in water method. The degree of caducity (high or low detachment of sporangium) was stated as a variable

character since it can be affected by the medium and method of sporangium production, while the pedicel length was a stable character since neither the medium nor the sporangium production has any effect on the sporangia with short and intermediate pedicel length (Al-Hedaithy and Tsao 1979b). The result of the present study differs from the earlier reports by the fact that cardamom isolates being intermediate in pedicel length, the pedicel length showed variation in solid and water culture and it was found to be directly proportional to caducity. The percentage of detached sporangia varied in solid and water culture. The description was comparable to those described for M isolates (*P. meadii*) of rubber and *Pricptiaral* var. *parasitice* isolates of *Cordyline terminalis* isolates of (Trichilo and Aragaki, 1982). Similarly analogous radial projections as outgrowths of the wall occur as a unique character on sporangia of many isolates of *P. nicotianae* var. *nicotianae* (Waterhouse *et al.*, 1983). These characters exhibited by some of the isolates of cardamom confirmed the presence of *P. nicotianae* var. *nicotianae* among cardamom isolates. Swelling along the sporangiophore was also characteristic of *P. nicotianae* var. *nicotianae* whereas swelling at the point of branching was characteristic of *P. meadii*, *P. citrophthora*, *P. citricola*, *P. colocasiae*, *P. infestans* and *P. phaseoli*.

The presence or absence of chlamydo spore is a significant parameter separating species such as *P. palmivora* from *P. megakarya* and *P. meadii* (Waterhouse *et al.*, 1983). None of the M isolates of rubber produced chlamydo spores on LBA (Lima Bean Agar medium), even after three months of incubation in natural light and dark at 20-25°C supporting the present study. Kunhimotto *et al* (1976) observed chlamydo spores in *P. nicotianae* var. *parasitica* in water when washed mycelial mats were kept in sterile water for 4 weeks at 16°C. M isolates of rubber produced chlamydo spores. Liyange and Wheeler (1991) reported that *P. meadii* isolates of Sri Lankan soils produced chlamydo spores in culture and they survived up to 12 weeks. In the present study also chlamydo spores were often noticed in Azhukal affected capsules when stored for long under low temperature (10°C) and also under ordinary conditions (28-32°C) indicated the association of a chlamydo spore forming species with Azhukal disease of cardamom and are considered as survival structures.

In the present investigation except one isolate, all the other isolates were found to be of A2 mating type thereby indicating the predominance of A2 mating type in cardamom. Some of the isolates were found to be bisexual /heterothallic producing sex organs in single culture. The

Table 6. Characters of *P. meadii* and *P. nicotianae* var. *nicotianae*

| A. Sporangial | As per the key (Stamps <i>et al.</i> , 1990) <i>P. meadii</i> | <i>P. nicotianae</i> var. <i>nicotianae</i> | Isolates studied <i>P. meadii</i> | <i>P. nicotianae</i> var. <i>nicotianae</i> |
|---------------------------------|---|--|--------------------------------------|--|
| Papillae pore narrow | + | Oc | + | + |
| > 1 apex | + | Oc | Oc | Oc |
| Caducous, pedicel length | M | Oc | M | M |
| Av./b ratio >1.6 | | * | >1.6 | >1.6 |
| ± spherical | * | * | | |
| Ellipsoidal | * | | | |
| Obpyriform | * | * | + | + |
| Obturbinate | | * | | |
| Elongated neck in water | + | | | |
| Distorted shapes | * | + | + | + |
| Hyphal projections Oc | | +/- | + | + |
| Lateral attachment | * | * | + | + |
| Intercalary | | + | | + |
| 45-75µm | * | + | + | + |
| Residual globule | + | | + | + |
| B. Sporangiphore | | | | |
| Aerial on agar | * | + | + | + |
| Basal swellings Oc. | + | | +Oc | |
| | Intercalary swellings | + | +Oc | |
| C. Antheridia | | | | |
| Amphigynous | * | + | + | + |
| ±spherical | * | + | + | + |
| Ellipsoidal or ovoid | * | | + | + |
| Stalk often eccentric | | + | | + |
| <30µm | * | + | + | + |
| 30-40µm | * | + | + | + |
| 40-50 µm | * | + | + | + |
| Wall rough with age | + | | + | + |
| D. Oospores | | | | |
| formed readily in the host | +/- | | | |
| Frequent on isolation | +/- | +/- | | |
| Often best in dual culture | + | + | + | + |
| <20-30m | * | + | + | + |
| 20- 30µm | * | + | + | + |
| 30-50µm | * | + | + | + |
| Wall thick (rel. dia) | * | + | + | + |
| Markedly aplerotic | * | + | + | + |
| E. Chlamydospores | | | | |
| Abundant | + | | | + |
| In some isolates only | + | | | + |
| <25µm | + | + | | + |
| 25-35µm | + | + | + | + |
| >35µm | | + | | + |
| F. Hyphal swellings on agar | | + | Oc | +Oc |
| In water round or angular | | + | + | \ |
| In chains or clumps | | | + | + |
| Large spherical | | | + | + |
| Spherical with radiating hyphae | | Oc | + | + |
| Growth at > 35°C | | +Oc | | + |

*character present from original descriptions and figures .

+Oc character has particular importance +/- present in some isolates

± more or less < less than > more than Oc occasional M medium

heterothallic nature of *P. meadii* occurring both as A1 and A2 compatibility type was in contrast to the above with some isolates showing self-fertility and one isolate sterile in pairings. Self-fertility in aging cultures (Brasier 1975; Peries and Dantanarayana, 1965) and self sterility in pairings (Brasier 1972) was also adjudgeable in the present study. Sansome *et al* (1991) reported self sterile isolates of *P. meadii*, which she attributed as due to subculturing.

Hybridization occurred between some of the cardamom isolates and *P. capsici* as shown by the sex organ formation on a band in between the two mating type. This was supported by the work of Boccas (1981). He obtained single oospore cultures from inter specific crosses of *P. megakarya* x *P. parasitica*, *P. parasitica* x *P. palmivora*, *P. parasitica* x *P. capsici*, *P. capsici* x *P. palmivora* and *P. cambivora* x *P. cinnamomi*. The present study showed the interspecific crosses between *P. nicotianae* x *P. capsici*.

According to Ho (1981) pigment production was characteristic of species such as *P. cactorum*, *P. cryptoega*, *P. cambivora*, *P. capsici*, *P. citricola*, *P. citrophthora*, *P. drechsleri* var. *drechsleri*, *P. erithroseptica*, *P. palmivora* MF4, *P. parasitica* var. *parasitica*, *P. parasitica* var. *nicotianae* and *P. vignae*. Kennedy and Duncan (1995) used CHT for pigment production to compare papillate *Phytophthora* species from raspberry namely *P. cactorum*, *P. citricola* and *P. syringae* with isolates of *P. cactorum* from apple and strawberry and observed the production of dark pigment as compared to the usual yellow pigment by *P. cactorum*, whereas no pigment production was observed in *P. citricola* and *P. syringae*.

A clear-cut differentiation could be obtained using the temperature relationship. The isolates which produced chlamydospores in culture and those which showed pigmentation in casein hydrolysate tyrosine agar medium were found growing at >35°C, whereas in others little or no growth was observed at this temperature. Lin and Chang (1986) studied the fruit rot of Durian caused by *P. palmivora* and explained the failure of these isolates to grow around 36°C and distinguished them from *P. nicotianae* and *P. nicotianae* var. *parasitica* which could grow well at temp above 35°C. Since Tucker (1931) many keys have used growth at minimum, maximum and optimum temperature as a useful criteria for classification at species level (Waterhouse *et al.*, 1983). Among the 57 isolates of *P. nicotianae* var. *parasitica* studied by Tucker, 7 isolates grew well at 35°C. Minimum temperature of *P. meadii* was reported as 10°C and Optimum temperature for mycelial growth was 28-30°C (Waterhouse, 1974a).

Identification of *Phytophthora* isolates of cardamom based on the Tabular key of Newhook *et al.* 1978; Ho, 1981; Stamps *et al.*, 1990

All the 25 isolates of cardamom studied were papillate with intermediate pedicel length (5-20 µm). Ho (1981) included species such as *P. botryosa*, *P. colocasiae*, *P. illicis*, *P. meadii* and *P. megakarya* MF3 in the group having caducous sporangia which shed in water with pedicel length 5-20 µm. *P. parasitica* var. *nicotianae* or *P. nicotianae* var. *nicotianae* was included in the non-caducous group with pedicel absent or non-uniform type. Among the caducous group with pedicel 5-20 µm, *P. colocasiae*, *P. illicis* were semi papillate type and hence the possibilities of their occurrence among cardamom isolates were ruled out. Internal proliferation is not observed in any of the cardamom isolates. The l/b ratio of the isolates were >1.6 with ovoid or ellipsoidal or obpyriform sporangia. Sex organs were produced readily and abundantly in intra and inter-specific crossings with distinct A1 and A2 compatibility types. Antheridia is entirely amphigynous, 2-20 µm long. Ability to produce chlamydospores varies with the isolates. In the first set chlamydospores were rare or absent and in the third set it is present at all temperatures. The first set included *P. botryosa* and *P. megakarya* whereas the second set included *P. meadii*, *P. nicotianae* var. *nicotianae* came under the third set due to the abundance of chlamydospores in culture. Thus based on the tabular key to the species of *Phytophthora* de Bary (Newhook *et al.*, 1978; Ho, 1981 and Stamps *et al.*, 1990) the isolates of cardamom were grouped into two categories. Category I having pigmentation in casein hydrolysate tyrosine agar with chlamydospore production and the ability to grow at >35°C and Category II having no pigmentation in casein hydrolysate tyrosine agar without chlamydospore production and no growth at 35°C. The pedicel length of the two groups fell in the intermediate group of 5- 20 µm. The above characters along with the other diagnostic characters in the key such as caducity, sporangial formation, nature of sex organs, hyphal swellings etc. all agreed with that of *P. nicotianae* var. *nicotianae* in the first category and *P. meadii* in the second category. Except for these characters all other characters were similar to the two categories and it was difficult to distinguish *P. meadii* from *P. nicotianae* var. *nicotianae* by simple morphological characters. Hence from the results it was found that the cultures with papillate sporangia of intermediate pedicel, hyphal swellings along sporangiophore, amphigynous antheridia spherical oogonia with chlamydospore production in culture and the ability to grow at >35°C along with

pigmentation in CHT medium were *P. nicotianae* var. *nicotianae* whereas the characters as above minus chlamyospore production, pigmentation and ability to grow at $>35^{\circ}\text{C}$ were *P. meadii*.

Characterization of the species

Based on the studies, two basic groups can be recognized in initiating capsule rot and leaf blight disease of cardamom. Group I having pigmentation in CHT whereas Group II without. Each group could again be divided into two depending on the formation of chlamyospores. Group I was found to be growing at high temp ($>35^{\circ}\text{C}$) whereas Group II has no growth at that temperature. According to the key of Ho (1981) the Group I having pigmentation, chlamyospore formation and growth at $>35^{\circ}\text{C}$ with occasional hyphal swellings from *P. parasitica* group is *P. nicotianae* var. *nicotianae*. The non--chlamyospore forming isolates without pigmentation and no growth at $>35^{\circ}\text{C}$ were invariably *P. meadii*. Moreover both of these groups readily produced sex organs with known A1 isolate of black pepper thereby establishing the predominance of A2 mating type of *P. meadii* and *P. nicotianae* in cardamom. Earlier only A1 isolate of *P. meadii* was reported (Sastry, 1989) Liyanage and Wheeler, 1989, and Rajalekshmi *et al.*, 1985). Hence in the light of the above studies it is confirmed that *P. Mc Rae* of A2 mating type is the predominant *Phytophthora* species involved in capsule rot and leaf blight occurring during the post monsoon period is due to *P. nicotianae* var. *nicotianae*. However there is certain amount of overlapping. *P. nicotianae* var. *nicotianae* can also cause capsule rot. As the pedicel length of none of the isolates fall in the range of $<5\mu\text{m}$, the possibility of the occurrence of *P. palmivora* can be ruled out.

Acknowledgements

The first author is grateful to Spices Board, Kochi for sanctioning study leave and also to the Director and staff of IISR Calicut for providing all the facilities for carrying out the research work.

References

- Al-Hedaithy, S.S.A. and Tsao, P.H. 1979(a). The effect of culture media and sporulation methods on caducity and pedicel length of sporangia in selected species of *Phytophthora*. *Mycologia*, 71: 392-401.
- Al-Hedaithy, S.S.A. and Tsao, P.H. 1979(b). Sporangium pedicel length in *Phytophthora* species and the consideration of its uniformity in determining sporangium caducity. *Trans. Brit. Mycol. Soc.* 72: 113.
- Anonymous. 1986. ICRI Annual Report. Myladumpara, pp.51.
- Boccas, B.R. 1981. Interspecific crosses between closely related heterothallic *Phytophthora* species. *Phytopathology*, 71: 60-65.
- Brassier, C.M. 1972. Observations on the sexual mechanism in *Phytophthora palmivora* and related species. *Trans. Brit. Mycol. Soc.* 58: 237-251.
- Brassier, C.M. 1975. Stimulation of sex organ formation in *Phytophthora* by antagonistic species of *Trichoderma* I. The effect in vitro. *New Phytol.* 74: 183-194.
- Brassier, C.M. and Griffin M.J. 1979. Taxonomy of *Phytophthora palmivora* of Cocoa. *Trans. Brit. Mycol. Soc.* 72: 111-143.
- Brassier, C.M., Hamm, P.B. and Hansen, E.M. 1993. Cultural characters, protein pattern and unusual mating behavior of *Phytophthora gonapodydes* isolates from Britain and North America. *Mycol. Res.* 97: 1287-1298.
- Chattopadhyaya, B., 1967. Diseases of plants yielding drugs, dyes and spices. Indian Council of Agricultural Research, New Delhi, p 100.
- Dantanarayana, D.M. Peries, O.S. and Des Liyanage A. 1984. Taxonomy of *Phytophthora* species isolated from Rubber in Sri Lanka. *Trans. Brit. Mycol. Soc.* 82: 113-126.
- Garrettson-Cornell, L., 1989. Technical papers 45.
- Ho, H.H. 1981. Synoptic key to the species of *Phytophthora*. *Mycologia*. 73(4): 705-713.
- Kennedy, D.M. and Duncan J.M. 1995. A papillate *Phytophthora* species with specificity to *Rubus*. *Mycol. Res.* 99: 57-68.
- Kunhimotto, R.K., Aragaki, M., Hunter, J.F. and Ko, W.H. 1976. *Phytophthora capsici*, corrected name for the cause of *Phytophthora* blight of *Macademia* recemes. *Phytopathology* 66: 546-548.
- Lim, T.K. and Chan, L.G. 1986. Fruit rot of durian caused by *phytophthora palmivora*. *Pertanika* 9: 269-276.
- Liyanage, A. Des., Peries, O.S., Dharmaratne, A. and Dantanarayana, D.M. 1983. Effect of weather factors on disease establishment and sporulation of *P. meadii* on Rubber pods. *J. of The Rubber Res. Inst. of Sri Lanka* 61: 41-48.
- Liyanage, N.I.S., Wheeler, B.E.J., 1989. Comparative morphology of *Phytophthora* species on rubber. *Plant Pathology*. 38: 592-597.
- Liyanage, N.I.S. and Wheeler, B.E.J. 1991. Survival of *Phytophthora meadii* on Sri Lankan soils. *Plant pathology* 40: 436-444.
- Menon, M.R. Sajoo, B.V., Ramakrishnan, C.K. and Remadevi, L. 1972. A new *Phytophthora* disease of cardamom (*Elettaria cardamomum*: L. Maton). *Curr. Sci.* 41: 231.
- Nair, R.R. and Menon, M.R. 1980. Azhukal disease of cardamom. In: *Proceedings of the workshop on Phytophthora disease of Tropical cultivated plants*. CPCRI, Kasaragod, pp 24-33.
- Newhook, F.J., Waterhouse, G.M. and Stamps, D.J. 1978. Tabular key to the species of *Phytophthora* de Bary. *Mycological papers*. 143.
- Peries, O.S. and Dantanarayana, D.M. 1965. Compatibility and variation in *Phytophthora* cultures isolated from *Hevea brasiliensis* in Ceylon. *Trans. Brit. Mycol. Soc.* 48: 631-637.
- Radha, K. and Joseph, T. 1974. Investigations on the bud rot (*Phytophthora palmivora*) of coconut PI. 480. Final Report. CPCRI, Kayamkulam pp.30.
- Rajalekshmi, V.K. and Annakkutty Joseph 1986. Production of sporangia by *Phytophthora meadii*. *Indian Phytopathology*. 39: 470-471.

- Rajalekshmi, V.K., Amakkutty Joseph and Arthasserry, S. 1985 Occurrence of two mating groups in *Phytophthora meadii* causing abnormal leaf fall disease of rubber in South India. *Trans. Brit. Mycol. Soc.* **85**: 723-726.
- Ramakrishnan, T.S. 1949. The occurrence of *Pythium vexans* de Bary in south India. *Indian Phytopath.* **2**: 27-30.
- Ribeiro, O.K., Erwin, D.C. Zentmyer, G.A., 1975. An improved synthetic medium for oospore production and germination of several *Phytophthora* species. *Mycologia* **67**: 101-1019.
- Sasthy, M.N.L. 1982. *Studies on species of Phytophthora affecting Plantation crops in Karnataka with special reference to Koleroga of arecanut and Black pepper*. Ph.D thesis, Dept. of Plant Pathology, College of Agriculture, Dharwad University, of Agri. Sciences, Bangalore.
- Sastry, M.N.L., Hegde, R.K. 1989. Variability of *Phytophthora* species obtained from plantations of Karnataka. *Indian Phytopath.* **42**: 421-425.
- Sansome, E.R., Brasier, C.M. and Harms, P.B., 1991. *Phytophthora meadii* may be a species hybrid. *Mycol. Res.* **95**: 273-277.
- Stamps, D.J., Waterhouse, O.M., Newhook, F.J. and Hall, O.S. 1990. Revised tabular key to the species of *Phytophthora* (Second edition). Mycological paper. 162 p.
- Timmer, L.W., Erwin, D.C. and Mc Cormic, W.H. 1970 Glucose inhibition and inheritance of pigment producing ability in *Phytophthora*. *Mycologia* **62**: 967-977.
- Thankamma, L., George M.K. and George K.V. 1968. Occurrence of two species of *Phytophthora* on *Hevea brasiliensis* in India. Rubber Board Bulletin 10, 33.
- Thankamma, L. and Pillai, P.N.R., 1973. Fruit rot and leaf rot disease of cardamom in India. *Plant Dis. Repr.* **58**: 767-769.
- Thomas, K. M. 1939. Report of sub-ordinate officers. Dept of Agric. 1938-1939. 130 p.
- Trichilo, P.J. and Aragaki, M. 1982 Sporangia! caducity and pedicel length of *P. nicotianae* var. *parasitica*. *Mycologia* **74**: 927-931.
- Tsao, P.H., Guy, S.O. 1977. Inhibition of *Mortierella* and *Pythium* in a *Phytophthora* isolation medium containing Hymexazole. *Phytopathology* **67**: 796-801.
- Tsao, P.H., Sarma, Y.R., Kasim, R., Mustika and Kueh, T.K. 1985 Variation in *Phytophthora palmivora* MF4 (*P. capsici*) isolates from black pepper in Indonesia and Malaya. *Phytopathology* **75**, 1315 (Abstract).
- Tucker, C.M. 1931. Taxonomy of the Genus *Phytophthora* De Bary Res. Bull: No. Agric. Expt. Stat. **153**: 1-208.
- Waterhouse G .M. 1961. Key to the species of *Phytophthora* de Bary. *Mycological paper*. No. 92 pp. 22.
- Waterhouse. G. M. 1974 (a) *Phytophthora palmivora* and some related species. In *Phytophthora* diseases of Cocoa. P.H. Gregory, ed. Longman, pp 51-70.
- Waterhouse, G.M., Newhook, F.J and Stamps, D.J. 1983. Present criteria in the classification of *Phytophthora*. In: *Phytophthora its Biology, Taxonomy, Ecology and Pathology*, (Ed) Erwin, D.C., Bartinicki-Garcia, S. and Tsao, P.H.; pp 139-147.
- Zentmyer, G.A., Kaosiri, T. and Idosu, G. 1977 Taxonomic variants in the *Phytophthora palmivora* complex. *Trans. Brit. Mycol Soc.* **69**: 329-332.