



## Molecular characterization of *Phytophthora nicotianae* associated with diseases of horticultural crops by RFLP of PCR internal transcribed spacer region of ribosomal DNA and AFLP fingerprints.

P. Chowdappa<sup>1</sup>

Central Plantation Crops Research Institute  
Regional Station, Vittal- 574 243, India  
E-mail: pallem @iihr.ernet.in

### ABSTRACT

Molecular characterization of *Phytophthora* isolates from tobacco, tomato, carnation, gerbera, crossandra and periwinkle was carried out using internal transcribed spacer(ITS) regions of rDNA gene repeat and amplified fragment length polymorphism (AFLP). All the isolates of *Phytophthora* were identical in morphology and ITS-RFLP patterns, indicating *P. nicotianae* and *P. parasitica* are synonymous. However, based on AFLP, four sub groups were evident within population of *P. nicotianae* (*P. parasitica*): Group I, includes isolates from tomato, tobacco and periwinkle. Isolates from carnation represents Group II. Isolates from gerbera falls under group III. The crossandra isolates formed a Group IV. Thus, present study showed the existence of four molecular sub groups within population of *P. nicotianae* (*P. parasitica*) in India.

**Key words:** *Phytophthora nicotianae*, AFLP, ITS-RFLP

### INTRODUCTION

*Phytophthora parasitica* (sensu Dastur, 1913) and *P. nicotianae* (sensu Waterhouse, 1963), are pathogenic to a wide range of hosts ranging from tree crops to herbaceous plants (Oudemans and Coffey, 1991). *Phytophthora nicotianae* has been reported as causal agent of diseases of tobacco, crossandra, carnation, gerbera and periwinkle and *P. parasitica* on tomato in India (Mehrotra and Aggarwal, 2001). These species were mainly identified based on morphological features (Stamps *et al*, 1990; Sarma *et al*, 2002). Morphological criteria have limitations in understanding genetic diversity and genetic distance between isolates of *P. nicotianae* and *P. parasitica*. Protein patterns (Erselius and de Vallavielle, 1984), serological analysis (Morton and Dukes, 1967), DNA - RFLP patterns (Forster and Coffey, 1991) and isozyme patterns (Oudemans and Coffey, 1991) were found to be useful in understanding genetic divergence between isolates belonging to *P. nicotianae* and *P. parasitica*. Endonuclease restriction digestion analysis of the ITS region of ribosomal DNA (ITS-RFLP) (Cooke and Duncan, 1997; Crawford *et al*, 1996; Chowdappa *et al*, 2003a, b, c; Chowdappa and Sarma, 2003) and AFLP finger prints were successful to separate species and also differences. The objective of the present study is to understand molecular diversity of *Phytophthora* isolates from tobacco, tomato carnation, gerbera, crossandra

and periwinkle in India based on ITS-RFLP and AFLP fingerprinting.

### MATERIAL AND METHODS

#### Fungal isolates

*Phytophthora nicotianae* isolates from tobacco, tomato, periwinkle, gerbera (05 isolates each), carnation and crossandra (04 isolates each) collected from Bangalore, Mysore, Tumkur and Hassan districts of Karnataka were used in this study. Reference cultures of *P. nicotianae* (IMI 3406160 and IMI 235604) were also included for comparison.

#### Morphological studies

Colony morphology of isolates, on carrot agar medium (CA) after three days of incubation in the dark at  $24 \pm 1^\circ\text{C}$ , was studied. Sporangia were produced on CA using mycelium-agar-disk-in-water method (Al-Hedaithy and Tsao, 1979). Observations were made on length and breadth of sporangia, presence of papilla, caducity and shapes. Observations on chlamydospore production were made on CA following incubation at  $18^\circ\text{C}$  in the dark. The mating types of isolates were determined by taking 5 mm dia. discs from advancing margins of 3-day-old cultures of opposite mating types and placed them 20 mm apart on CA. The inoculated plates were incubated

<sup>1</sup>Present address: Division of Plant Pathology, Indian Institute of Horticultural Research, Bangalore – 560 089, India

at 20°C in the dark for 15-20 days. The influence of temperature on vegetative growth on CA was determined in the dark at temperature ranging from 6 to 39°C with an increment of 3°C.

### Molecular studies

The fungal mycelium was grown in 100 ml of V8 juice in a 250 ml flask for 3 days at 25°C on orbital shaker at 100 rpm. DNA was extracted from freeze dried vegetative mycelium following a method of Raeder and Broda (1985). PCR amplification of the ITS region was performed using with universal primer pair ITS1 and ITS 4 (White *et al*, 1990). Each 50 µl reaction mixture contained 100 ng of DNA, 50 pmol of each primer, 200 µM each dNTP, 5 µl of 10 x PCR buffer and 10 units of Taq DNA polymerase enzyme. PCR was carried out with following cycling parameters. A 4 minute denaturation at 94°C, followed by 34 cycles of 60s at 94°C, 60s at 55°C, and 1.5 min at 72°C and then a 5 min extension at 72°C. PCR products (5µl) were electrophoresed in a 2% agarose gel with Tris-Borate-EDTA (TBE) buffer, stained with ethidium bromide and visualised under UV illumination. PCR product was digested with restriction enzymes namely *Hinf* I, *Msp* I, *Alu* I, *Hae* III, *Rsa* I, *Taq* I and *EcoR* V. Restriction digestions were performed in 10 µl reaction containing 5 µl PCR product, 1 µl 10 x restriction buffer, 3.6 µl PCR grade water, 0.1 µl Bovine serum albumin and restriction enzyme (3 U/reaction) at 37°C for overnight (16h). Digestion products were electrophoresed in 2.5% LE agarose. The size of restriction fragments was determined by comparison of fragments migration with that of known marker fragments (100bp molecular size ladder).

Amplified fragment length polymorphism (AFLP) analysis was carried out as per the method of Muller *et al* (1996). Fungal genomic DNA (500 ng) was digested by *Pst* (20 units), ligated to *Pst* adaptor at 37°C for 4h and pre-amplification was carried out at 94°C for 4 min for initial denaturation followed by 34 cycles of denaturation for 1 min at 94°C, annealing for 1 min at 60°C, extension for 1.5 min at 72°C and a final extension step of 5min at 72°C. Pre-amplified PCR products were then diluted 1:100 in TE buffer and amplified with five AFLP primers, A(5'-GACTGCGTACATGCAGGT3', B(5'-GACTGCGTACATGCAGGA3'), C, 5'-GACTGCGTACATGCAGGC3'), D (5'-GACTGCGTACATGCAGAC3'), E(5'-GACTGCGTACATGCAGCG-3') each with two selective bases at its 3' end underlined, adopting above thermal cycling parameters. The PCR products were electrophoresed in 2% agarose gels in Tris-Borate-EDTA (TBE) buffer and visualized under UV after staining with ethidium bromide


**Table 1. Sporangial measurement in *P. nicotianae* isolates**

Isolate No	Length*(µm)	Breadth (µm)	L/B ratio(µm)	Mating type
Tobacco				
OH 37	41.50 ± 1.21	29.50 ± 0.77	1.40 ± 0.77	A2
OH 38	43.07 ± 0.93	31.92 ± 0.61	1.33 ± 0.02	A2
OH 39	41.00 ± 1.08	30.20 ± 0.71	1.37 ± 0.02	A2
OH 40	36.70 ± 1.21	24.92 ± 0.54	1.47 ± 0.36	A2
OH 41	40.93 ± 1.91	27.80 ± 0.69	1.41 ± 0.04	A2
Tomato				
OH 42	38.50 ± 1.67	28.50 ± 1.14	1.30 ± 0.06	A2
OH 43	42.50 ± 0.86	32.25 ± 0.74	1.30 ± 0.02	A2
OH 44	40.00 ± 1.30	29.57 ± 0.83	1.37 ± 0.05	A2
OH 45	40.50 ± 0.77	29.00 ± 0.63	1.39 ± 0.03	A2
OH 46	39.00 ± 1.71	25.50 ± 1.00	1.38 ± 0.08	A2
Periwinkle				
OH 47	41.66 ± 1.45	28.54 ± 1.23	1.43 ± 0.03	A2
OH 48	42.75 ± 1.14	30.25 ± 0.96	1.41 ± 0.06	A2
OH 49	41.25 ± 1.53	28.75 ± 0.78	1.43 ± 0.02	A2
OH 50	41.66 ± 1.63	30.41 ± 1.09	1.36 ± 0.03	A2
OH 51	39.58 ± 1.81	27.91 ± 1.37	1.41 ± 0.06	A2
Carnation				
OH 52	38.61 ± 1.52	27.22 ± 1.32	1.41 ± 0.02	A2
OH 53	38.50 ± 1.67	28.50 ± 1.14	1.30 ± 0.06	A2
OH 54	44.44 ± 1.39	30.55 ± 1.51	1.46 ± 0.06	A2
OH 55	38.75 ± 1.39	27.25 ± 1.20	1.45 ± 0.48	A2
OH 56	46.50 ± 2.74	31.50 ± 1.53	1.46 ± 0.03	A2
Crossandra				
OH 57	43.25 ± 1.17	28.75 ± 1.28	1.53 ± 0.06	A2
OH 58	45.00 ± 1.32	31.11 ± 1.42	1.46 ± 0.05	A2
OH 59	40.00 ± 1.23	27.00 ± 0.93	1.48 ± 0.04	A2
OH 60	46.38 ± 2.28	30.31 ± 1.34	1.53 ± 0.05	A2
OH 61	40.55 ± 1.70	27.77 ± 1.32	1.46 ± 0.05	A2
Gerbera				
OH 62	45.50 ± 1.88	30.25 ± 1.24	1.49 ± 0.04	A2
OH 63	41.66 ± 1.63	30.41 ± 1.09	1.36 ± 0.03	A2
OH64	41.25 ± 1.53	28.75 ± 0.78	1.43 ± 0.02	A2
OH 65	44.07 ± 1.00	30.47 ± 0.90	1.44 ± 0.04	A2
OH 66	42.91 ± 1.49	28.33 ± 1.27	1.51 ± 0.05	A2

\*Mean of 100 sporangia; ± standard deviation

## RESULTS AND DISCUSSION

*Phytophthora nicotianae* isolates isolated from tobacco, tomato, periwinkle, carnation, crossandra and gerbera displayed no distinctive colony morphology (dense whitish aerial mycelium). Most isolates produced spherical, ovoid, ellipsoid and persistent sporangia on irregular / loose sympodial sporangiophore (Fig1). Sporangial sizes ranged from 36.70 to 46.50 x 24.92 to 32.25 µm (Table1). L:B ratios of sporangia ranged from 1.31 to 1.53. Chlamydospores were produced readily by all isolates on CA incubated in the dark at 18°C. Chlamydospores were single, spherical, terminal or intercalary and pale yellow in colour (Fig 1). The mean chlamydospores diameter ranged from 20 to 30 µm (Table 2). None of the isolates produced sexual structures. When paired with isolates of *P. capsici* of known mating types, all isolates produced sexual



**Fig 1. Morphology of *P. nicotianae* isolates, Top left, sporangia, Top right, chlamydospores, Bottom left, oogonia, Bottom right, oospores**

structures (Fig 1). Size of oogonia ranged from 26 to 29  $\mu\text{m}$  and oospores varied from 24 to 27  $\mu\text{m}$  (Table 2). Minimum temperature for growth of all isolates was 9°C; optimum was 24 - 30°C; and maximum was 33°C.


Thus, the present study showed that sporangial, chlamydospore, oogonial and oospore characteristics of *P. nicotianae* isolates from periwinkle, carnation, gerbera, tobacco and crossandra are similar to those of *P. parasitica* isolates from tomato. Waterhouse (1963) created two morphological varieties; *P. nicotianae* var. *nicotianae* and *P. nicotianae* var. *parasitica*, which were based on Ashby (1928) microspore and macrospora, respectively. Further morphological studies of Ho and Jong (1989) in isolates from tobacco and other hosts clearly showed that var. *nicotianae* and var. *parasitica* could not be differentiated. Separation of the varieties *nicotianae* and *parasitica* based on morphological criteria was also questioned (Gallegly, 1983). Oudemans and Coffey (1991) and Hall (1993) reported that oogonia, oospore and sporangial dimensions are similar among two groups.

When DNA was amplified with universal primer pair (ITS1 and ITS4), there was single band of 920 bp (Fig 2) for all isolates of *P. nicotianae* and reference cultures of *P. nicotianae* (IMI 3406160 and IMI 235604). All isolates had identical ITS - RFLP patterns when PCR product was digested with enzymes *Hinf I*, *Msp I*, *Alu I*, *Hae III* and *Rsa I* (Fig 3). Restriction fragment sizes of the ITS region of rDNA digested with different enzymes are also identical.

**Table 2. Dimensions of chlamydospores, oogonia and oospores of *P. nicotianae* isolates**

Isolate	Chlamydospores ( $\mu\text{m}$ )	Oogonia ( $\mu\text{m}$ )	Oospores ( $\mu\text{m}$ )
Tobacco			
OH 37	24.38 $\pm$ 1.75*	28.34 $\pm$ 2.10	25.90 $\pm$ 2.00
OH 38	22.78 $\pm$ 2.30	27.26 $\pm$ 1.90	26.24 $\pm$ 1.75
OH 39	28.20 $\pm$ 2.60	28.75 $\pm$ 2.00	25.34 $\pm$ 2.30
OH 40	23.34 $\pm$ 3.24	25.20 $\pm$ 1.95	24.37 $\pm$ 2.30
OH 41	23.28 $\pm$ 1.95	27.32 $\pm$ 2.60	25.20 $\pm$ 2.60
Tomato			
OH 42	27.34 $\pm$ 2.02	29.75 $\pm$ 2.72	26.37 $\pm$ 2.27
OH 43	28.28 $\pm$ 2.78	27.14 $\pm$ 2.02	25.40 $\pm$ 2.50
OH 44	23.38 $\pm$ 2.50	27.30 $\pm$ 2.50	26.32 $\pm$ 2.71
OH 45	26.72 $\pm$ 2.31	28.14 $\pm$ 2.60	26.37 $\pm$ 2.80
OH 46	24.32 $\pm$ 1.31	27.20 $\pm$ 1.90	25.27 $\pm$ 1.70
Periwinkle			
OH 47	20.24 $\pm$ 1.32	28.14 $\pm$ 2.60	25.32 $\pm$ 2.75
OH 48	26.32 $\pm$ 2.34	28.20 $\pm$ 2.80	26.97 $\pm$ 2.60
OH 49	26.78 $\pm$ 2.14	27.20 $\pm$ 1.97	24.32 $\pm$ 2.32
OH 50	28.24 $\pm$ 3.21	27.30 $\pm$ 2.30	25.84 $\pm$ 2.00
OH 51	24.32 $\pm$ 2.20	27.10 $\pm$ 2.02	26.80 $\pm$ 2.72
Carnation			
OH 52	28.45 $\pm$ 2.10	28.32 $\pm$ 2.00	27.40 $\pm$ 1.72
OH 53	30.25 $\pm$ 1.78	27.20 $\pm$ 1.80	26.68 $\pm$ 2.34
OH 54	28.75 $\pm$ 2.24	27.45 $\pm$ 2.20	27.35 $\pm$ 2.40
OH 55	25.62 $\pm$ 3.14	28.47 $\pm$ 2.15	26.40 $\pm$ 1.80
OH 56	25.34 $\pm$ 2.45	27.90 $\pm$ 2.70	25.21 $\pm$ 2.30
Crossandra			
OH 57	26.25 $\pm$ 2.41	29.30 $\pm$ 1.60	24.00 $\pm$ 2.26
OH 58	20.35 $\pm$ 2.35	26.40 $\pm$ 2.30	25.34 $\pm$ 1.60
OH 59	20.75 $\pm$ 2.48	25.30 $\pm$ 1.60	26.73 $\pm$ 1.70
OH 60	24.34 $\pm$ 1.82	28.17 $\pm$ 1.20	27.40 $\pm$ 1.20
OH 61	25.75 $\pm$ 2.35	26.30 $\pm$ 2.00	24.37 $\pm$ 1.78
Gerbera			
OH 62	27.32 $\pm$ 1.32	27.34 $\pm$ 2.06	26.75 $\pm$ 1.80
OH 63	26.31 $\pm$ 2.71	26.48 $\pm$ 2.68	25.30 $\pm$ 2.07
OH 64	28.25 $\pm$ 2.24	27.84 $\pm$ 2.50	25.24 $\pm$ 2.01
OH 65	26.32 $\pm$ 2.18	26.34 $\pm$ 1.85	25.70 $\pm$ 2.07
OH 66	22.35 $\pm$ 2.14	27.84 $\pm$ 2.00	26.87 $\pm$ 2.80

\*Mean of 100 reproductive structures  $\pm$  standard deviation



**Fig 2. PCR amplification of ITS region of rDNA Lane1, Marker, 100bp ladder, Lane2, Periwinkle, Lane 3, gerbera, Lane,4,carnation, Lane,5, crossandra**




Fig 3. ITS-RFLP patterns generated with *Hinf I* (Lane 2-5) and *TaqI* (Lane 6-9). Lane 1 and 10, Marker (100bp) Lane 2 and 6, periwinkle, Lane 3 and 7, gerbera, Lane 4 and 8, carnation, Lane 5 and 9, crossandra

Protein profiles (Erselius and de Vallavielle, 1984), serological reactions (Morton and Dukes, 1967), RFLP-DNA patterns (Forster and Coffey, 1991) of *P. nicotianae* isolates from tobacco were almost identical to those of *P. nicotianae* isolates from other crops. ITS-RFLP data obtained in the present study also showed that *P. nicotianae* isolates from tobacco, tomato, periwinkle, carnation, gerbera and crossandra were identical. As varieties *nicotianae* and *parasitica* are identical based on morphological and molecular criteria, many researchers proposed (Ho and Jang, 1989; Oudemans and Coffey, 1991; Hall, 1993) that the name *P. nicotianae* be eliminated and that all isolates from tobacco and other hosts should be merged under the name *P. parasitica*.

Among the isolates of *P. nicotianae*, four AFLP finger print groups were evident based on the combined results of five primers (A, B, C, D and E). The AFLP finger prints generated with primer E is shown in Fig 4. Group I includes isolates from tomato, tobacco and periwinkle. Isolates from carnation represents Group II. Isolates from gerbera falls under group III. The crossandra isolates formed a Group IV. Although there was no intra specific ITS variation among isolates of *P. nicotianae*, host specialized groups from tobacco, tomato, periwinkle, gerbera, carnation, and crossandra were found based on AFLP fingerprints.

There is considerable amount of evidence to show that some isolates of *P. nicotianae* are host-specific. For instance, isolates from okra and sesame were host specific, while isolates from *Peperomia obtusifolia* and *Saintpaulia ionatha* had different host ranges (Ho and Jong, 1989). Isolates from carnation and tomato were more virulent on




Fig 4. AFLP finger prints of *Phytophthora nicotianae* isolates Lane 1, bp ladder (size marker), Lane 2, periwinkle, Lane 3, gerbera, Lane 4, carnation, Lane 5, crossandra

carnation and tomato respectively than other hosts. Thus, the present study clearly indicated that *P. nicotianae* isolates from tobacco, periwinkle, carnation, gerbera and crossandra and *P. parasitica* isolates from tomato were identical. The AFLP finger prints were useful in identifying host specialized groups within *P. parasitica* (*P. nicotianae*). While screening the accessions for disease resistance, host specialized AFLP molecular groups should be used rather than morphologically defined isolates.

## ACKNOWLEDGEMENT

The author is grateful to the Indian Council of Agricultural Research, New Delhi for financial assistance in the form of National Network Project on *Phytophthora* diseases of horticultural crops. He is also thankful to Dr. N. Ramachandran, Head, Division of Plant Pathology, Indian Institute of Horticultural Research, Bangalore, for *Phytophthora* cultures from gerbera, carnation and crossandra, and, to Dr. M. M. Shenoi, Head, CTCRI Regional Station, Hunsur, for *Phytophthora* infected samples of tobacco.

## REFERENCES

- Al-Hedaithy, S. S. A. and Tsao, P. H. 1979. Sporangium pedicel length in *Phytophthora* species and the consideration of its uniformity in determining sporangium caducity. *Trans. Brit. Mycol. Soc.*, **72**: 1-13.
- Ashby, S. F. 1928. The oospores of *Phytophthora nicotianae* Br. de Haan with notes on the taxonomy of *P. parasitica* Dastur. *Trans. Brit. Mycol. Soc.*, **13**: 86-95

- Chowdappa, P., Brayford, D., Smith, J. and Flood, J. J. 2003a. Molecular discrimination of *Phytophthora* isolates on cocoa and their relationship with coconut, black pepper and bell pepper isolates based on rDNA repeat and AFLP finger prints. *Curr. Sci.*, **84**: 1235 – 1238.
- Chowdappa, P., Brayford, D., Smith, J. and Flood J. J. 2003b. Identification of *Phytophthora* species affecting plantation crops by RFLP of PCR amplified ITS region of ribosomal RNA. *Curr. Sci.*, **85**: 34-36.
- Chowdappa, P., Brayford, D., Smith, J. and Flood, J. J. 2003c. Identity of *Phytophthora* associated with arecanut and its relationship rubber and cardamom isolates based on RFLP of PCR- amplified ITS region of rDNA and AFLP fingerprints. *Curr.Sci.*, **85**: 585-587.
- Chowdappa, P. and Sarma, Y. R. 2003. Application of molecular markers for resolving taxonomic complexities in *Phytophthora*. *J. Plantn. Crops.*, **31**: 1-12
- Cooke, D. E. L. and Duncan, J. M. 1997. Phylogenetic analysis of *Phytophthora* species based on ITS1 and ITS2 sequences of the ribosomal RNA gene repeat. *Mycol.Res.*, **101**: 667-677
- Crowford, A. R., Bassam, B. J., Drenth, A., Maclean, D. J. and Irwin, J. A. G. 1996. Evolutionary relationships among *Phytophthora* species deduced from rDNA sequence analysis. *Mycol. Res.*, **100**: 437-443.
- Dastur, J. F. 1913. *Phytophthora parasitica* n.sp, a new disease of castor oil plant. *Mem. Dep. Agric. Bot. Ser.*, **5**: 177-231.
- Erselius, L. J. and de Vallavieille, C. 1984. Variation in protein profiles in *Phytophthora*: comparison of six species. *Trans. Brit. Mycol. Soc.*, **83**: 463-472
- Forster, H. and Coffey, M. D. 1991. Approaches to the taxonomy of *Phytophthora* using polymorphisms in mitochondrial and nuclear DNA. p164-183. In: *Phytophthora*. Leucas, J.A. Shattock, R.C. Shaw.D.S. and Cooke, L.R.(ed.) Cambridge Uni. Press, Cambridge, UK
- Gallegly, M. E. 1983. New criteria for classifying *Phytophthora* and critique of existing approaches.p167-172. In: *Phytophthora: its Biology, Taxonomy, Ecology and Pathology*. Erwin, D.C. Bartnicki-Garcia, S. and Tsao, P.H.(ed.) American Phytopathological Society, St. Paul.
- Hall, G. 1993. An integrated approach to the analysis of variation in *Phytophthora nicotianae* and a redescription of the species. *Mycol. Res.*, **97**: 559-574.
- Ho, H. H. and Jong, S. C. 1989. *Phytophthora nicotianae* (*P. parasitica*). *Mycotaxon*, **36**: 243-276.
- Mehrotra, R. S. and Aggarwal, A. 2001. *Phytophthora* diseases in India. SMPS, Dehradun, p 322
- Morton, D. J. and Dukes, P. D. 1967. Serological differentiation of *Pythium aphanidermatum* from *Phytophthora parasitica* var. *nicotianae* and *Phytophthora parasitica*. *Nature*, **213**: 923.
- Mueller, U. G., Lipari, S. E. and Milgroom, M. G. 1996. Amplified fragment length polymorphism finger printing of symbiotic fungi cultured by fungus-growing ant *Cyphomyrex minutus*. *Mol. Ecology*, **5**: 119-122.
- Oudemans, P. and Coffey, M. D. 1991. A revised systematics of twelve papillate *Phytophthora* species based on isozyme analysis. *Mycol. Res.*, **95**: 1025-1046
- Raeder, U. and Broda. P. 1985. Rapid preparation of DNA from filamentous fungi. *Lett. Appl. Microbiol.*, **1**: 17-20.
- Sarma, Y R., Chowdappa, P. and Anandaraj. M. 2002. *Phytophthora* disease of plantation crops in India.p149-187. In: IPM system in Agriculture: Key pathogens and diseases .Vol. 8. Upadhyay, R.K. Arora, D. K. and Dubey, O. P. (ed.). Aditya Books Pvt. Ltd., New Delhi
- Stamps, D. J., Waterhouse, G. M., Newhook, F. J and Hall, G. S. 1990. Revised tabular key to the species of the *Phytophthora*, *Mycological papers* 162, CAB International Mycological Institute, Kew, London, p 28
- Waterhouse, G. M. 1963. Key to the species of *Phytophthora* de Bary; *Mycological papers* (CMI), **92**: 1-22
- White, T. J., Bruns, T., Lee, S. and Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. P 315-322. In: PCR protocols- a guide to methods and applications .Innis, M A. Gelfand, D. H. Sninsky, J. J. and White, T. J(ed. ). Academic Press, San Diego.

(MS Received 3 November, 2006, Revised 2 June 2007)